

# Signal Reporter Assay Kit

## SRE Reporter Assay Kit

Cat. no. 336841 CCS-010L

### For study of pathway activity

The Signal SRE Reporter Assay Kit is designed to monitor the activity of Serum Response Factor (SRF)-mediated signal transduction pathways in cultured cells. The ternary complex factors, TCR and Elk-1, form a complex with the SRF over the serum response element (SRE), and activate gene expression. The Elk-1 protein is phosphorylated by mitogen-activated protein kinase (MAPK), causing increased DNA binding, ternary complex formation, and transcriptional activation of target genes. The SRE reporter is designed for measuring the activation of the SRF and the MAPK signal transduction pathway. The SRE reporter is a mixture of an inducible SRE-responsive luciferase construct and constitutively expressing Renilla construct (40:1). The SRE-responsive luciferase construct encodes the firefly luciferase reporter gene under the control of a minimal (m)CMV promoter and tandem repeats of the SRE. We have experimentally optimized the number of response elements as well as the intervening sequence between response elements to maximize the signal to noise ratio. The constitutively expressing Renilla construct encodes the Renilla luciferase reporter gene under the control of a CMV immediately early enhancer/promoter and acts as an internal control for normalizing transfection efficiencies and monitoring cell viability. Using a simple dual-luciferase assay, you can easily monitor the activity of SRF-mediated signaling pathways and determine the effect of various treatments, such as gene knockdown, over-expression, and chemical compounds on those pathways.

For further details, consult the *Signal Reporter Assay Handbook*.

### Shipping and storage

The Signal Reporter Assay Kit is shipped on dry ice or blue ice. Store at  $-20^{\circ}\text{C}$ . When stored under these conditions, this product can be used for 6 months from the date of receipt without reduction in performance.

**Note:** Open the package and store the products appropriately immediately on receipt.

### Performance data

Visit the Website to view performance data for this product.



---

Sample & Assay Technologies

## Contents

| Product                   | Contents                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                              | Cat. no. |
|---------------------------|-------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|----------|
| Signal Reporter Assay Kit | For 500 reporter assays, 500 negative control assays, and 250 positive control assays when using the recommended 96-well plate transfection protocol; 500 $\mu$ l (100 ng/ $\mu$ l) Reporter (40:1 mixture of inducible firefly luciferase construct and constitutively expressing Renilla luciferase construct); 500 $\mu$ l (100 ng/ $\mu$ l) Negative control (40:1 mixture of non-inducible firefly luciferase construct and constitutively expressing Renilla luciferase construct); 250 $\mu$ l (100 ng/ $\mu$ l) Positive control (40:1:1 mixture of constitutively expressing GFP, constitutively expressing firefly luciferase, and constitutively expressing Renilla luciferase constructs) | Varies   |

Signal Reporter Assay Kit products are intended for molecular biology applications. These products are not intended for the diagnosis, prevention, or treatment of a disease.

For up-to-date licensing information and product-specific disclaimers, see the respective QIAGEN kit handbook or user manual. QIAGEN kit handbooks and user manuals are available at [www.qiagen.com](http://www.qiagen.com) or can be requested from QIAGEN Technical Services or your local distributor.

Trademarks: QIAGEN® (QIAGEN Group).

1065997 03/2011 © 2011 QIAGEN, all rights reserved.

[www.qiagen.com](http://www.qiagen.com)

Australia ■ 1-800-243-800

Austria ■ 0800/281010

Belgium ■ 0800-79612

Brazil ■ 0800-557779

Canada ■ 800-572-9613

China ■ 8621-3865-3865

Denmark ■ 80-885945

Finland ■ 0800-914416

France ■ 01-60-920-930

Germany ■ 02103-29-12000

Hong Kong ■ 800 933 965

Ireland ■ 1800 555 049

Italy ■ 800-787980

Japan ■ 03-6890-7300

Korea (South) ■ 080-000-7145

Luxembourg ■ 8002 2076

Mexico ■ 01-800-7742-436

The Netherlands ■ 0800 0229592

Norway ■ 800-18859

Singapore ■ 1800-742-4368

Spain ■ 91-630-7050

Sweden ■ 020-790282

Switzerland ■ 055-254-22-11

UK ■ 01293-422-911

USA ■ 800-426-8157



Sample & Assay Technologies