

Signal Reporter Assay Kit

SMAD Reporter Assay Kit (GFP)

Cat. no. 336841 CCS-017G

For study of pathway activity

The Signal SMAD Reporter Assay Kit is designed to monitor the activity of the TGF β -induced signal transduction pathway in cultured cells. The transforming growth factor β (TGF β) signaling pathway is involved in many cellular processes, including cell cycle arrest, differentiation, homeostasis, and immunosuppression. TGF β signaling induces phosphorylation and activation of the SMAD2 and SMAD3 proteins, which then form complexes with the mediator SMAD4. These SMAD complexes then translocate to the nucleus, where they activate the expression of TGF β -responsive genes. The SMAD reporter expresses the Monster GFP reporter gene upon activation of TGF β -responsive cell signaling pathways. The SMAD-responsive GFP construct encodes the Monster GFP gene under the control of a minimal (m)CMV promoter and tandem repeats of the SMAD binding element (SBE). We have experimentally optimized the number of response elements as well as the intervening sequence between response elements to maximize the signal to noise ratio. Using a simple GFP assay, you can easily monitor the activity of SMAD-mediated signaling pathways and determine the effect of various treatments, such as gene knockdown, over-expression, and chemical compounds on those pathways.

For further details, consult the *Signal Reporter Assay Handbook*.

Shipping and storage

The Signal Reporter Assay Kit is shipped on dry ice or blue ice. Store at -20°C . When stored under these conditions, this product can be used for 6 months from the date of receipt without reduction in performance.

Note: Open the package and store the products appropriately immediately on receipt.

Performance data

Visit the Website to view performance data for this product.



Sample & Assay Technologies

Contents

Product	Contents	Cat. no.
Signal Reporter Assay Kit	For 500 reporter assays, 500 negative control assays, and 250 positive control assays when using the recommended 96-well plate transfection protocol; 500 μ l (100 ng/ μ l) Reporter (40:1 mixture of inducible firefly luciferase construct and constitutively expressing Renilla luciferase construct); 500 μ l (100 ng/ μ l) Negative control (40:1 mixture of non-inducible firefly luciferase construct and constitutively expressing Renilla luciferase construct); 250 μ l (100 ng/ μ l) Positive control (40:1:1 mixture of constitutively expressing GFP, constitutively expressing firefly luciferase, and constitutively expressing Renilla luciferase constructs)	Varies

Signal Reporter Assay Kit products are intended for molecular biology applications. These products are not intended for the diagnosis, prevention, or treatment of a disease.

For up-to-date licensing information and product-specific disclaimers, see the respective QIAGEN kit handbook or user manual. QIAGEN kit handbooks and user manuals are available at www.qiagen.com or can be requested from QIAGEN Technical Services or your local distributor.

Trademarks: QIAGEN® (QIAGEN Group).

1065997 03/2011 © 2011 QIAGEN, all rights reserved.

www.qiagen.com

Australia ■ 1-800-243-800

Austria ■ 0800/281010

Belgium ■ 0800-79612

Brazil ■ 0800-557779

Canada ■ 800-572-9613

China ■ 8621-3865-3865

Denmark ■ 80-885945

Finland ■ 0800-914416

France ■ 01-60-920-930

Germany ■ 02103-29-12000

Hong Kong ■ 800 933 965

Ireland ■ 1800 555 049

Italy ■ 800-787980

Japan ■ 03-6890-7300

Korea (South) ■ 080-000-7145

Luxembourg ■ 8002 2076

Mexico ■ 01-800-7742-436

The Netherlands ■ 0800 0229592

Norway ■ 800-18859

Singapore ■ 1800-742-4368

Spain ■ 91-630-7050

Sweden ■ 020-790282

Switzerland ■ 055-254-22-11

UK ■ 01293-422-911

USA ■ 800-426-8157



Sample & Assay Technologies