

Signal Reporter Assay Kit

FOXO Reporter Assay Kit

Cat. no. 336841 CCS-1022L

For study of pathway activity

The Signal FOXO Reporter Assay Kit is designed to measure the activity of PI3K/AKT signaling pathway via the transcriptional activity of the FOXO transcription factors, which are a subfamily of the Forkhead transcription factor family. The FOXO reporter is a mixture of a FOXO-responsive luciferase construct and a constitutively expressing Renilla element (40:1). The FOXO-responsive luciferase construct encodes the firefly luciferase reporter gene under the control of a minimal (m)CMV promoter and tandem repeats of the FOXO transcriptional response element. The number of response elements and the intervening sequence between these response elements has been experimentally optimized to maximize the signal to noise ratio. This construct monitors both increases and decreases in the transcriptional activity of FOXO, and hence the activity of the PI3K/AKT signaling pathway. The constitutively expressing Renilla construct encodes the Renilla luciferase reporter gene under the control of a CMV immediately early enhancer/promoter and acts as an internal control for normalizing transfection efficiencies and monitoring cell viability. Using a simple dual-luciferase assay, the activity of the PI3K/AKT signaling pathway can be easily measured in response to various treatments, such as gene knockdown, over-expression, and chemical compounds on those pathways.

For further details, consult the *Signal Reporter Assay Handbook*.

Shipping and storage

The Signal Reporter Assay Kit is shipped on dry ice or blue ice. Store at -20°C . When stored under these conditions, this product can be used for 6 months from the date of receipt without reduction in performance.

Note: Open the package and store the products appropriately immediately on receipt.

Performance data

Visit the Website to view performance data for this product.



Sample & Assay Technologies

Contents

Product	Contents	Cat. no.
Signal Reporter Assay Kit	For 500 reporter assays, 500 negative control assays, and 250 positive control assays when using the recommended 96-well plate transfection protocol; 500 μ l (100 ng/ μ l) Reporter (40:1 mixture of inducible firefly luciferase construct and constitutively expressing Renilla luciferase construct); 500 μ l (100 ng/ μ l) Negative control (40:1 mixture of non-inducible firefly luciferase construct and constitutively expressing Renilla luciferase construct); 250 μ l (100 ng/ μ l) Positive control (40:1:1 mixture of constitutively expressing GFP, constitutively expressing firefly luciferase, and constitutively expressing Renilla luciferase constructs)	Varies

Signal Reporter Assay Kit products are intended for molecular biology applications. These products are not intended for the diagnosis, prevention, or treatment of a disease.

For up-to-date licensing information and product-specific disclaimers, see the respective QIAGEN kit handbook or user manual. QIAGEN kit handbooks and user manuals are available at www.qiagen.com or can be requested from QIAGEN Technical Services or your local distributor.

Trademarks: QIAGEN® (QIAGEN Group).

1065997 03/2011 © 2011 QIAGEN, all rights reserved.

www.qiagen.com

Australia ■ 1-800-243-800

Austria ■ 0800/281010

Belgium ■ 0800-79612

Brazil ■ 0800-557779

Canada ■ 800-572-9613

China ■ 8621-3865-3865

Denmark ■ 80-885945

Finland ■ 0800-914416

France ■ 01-60-920-930

Germany ■ 02103-29-12000

Hong Kong ■ 800 933 965

Ireland ■ 1800 555 049

Italy ■ 800-787980

Japan ■ 03-6890-7300

Korea (South) ■ 080-000-7145

Luxembourg ■ 8002 2076

Mexico ■ 01-800-7742-436

The Netherlands ■ 0800 0229592

Norway ■ 800-18859

Singapore ■ 1800-742-4368

Spain ■ 91-630-7050

Sweden ■ 020-790282

Switzerland ■ 055-254-22-11

UK ■ 01293-422-911

USA ■ 800-426-8157



Sample & Assay Technologies