

Cignal Reporter Assay Kit

PPAR Reporter Assay Kit

Cat. no. 336841 CCS-3026L

For study of pathway activity

The Cignal PPAR Reporter Assay Kit is designed to measure the transcriptional activity of peroxisome proliferator-activated receptors (PPARs). Drugs targeting PPARs may serve as treatments for diseases associated with aging, such as arteriosclerosis and diabetes. The PPAR reporter is a mixture of a PPAR-responsive luciferase construct and a constitutively expressing Renilla element (40:1). The PPAR-responsive luciferase construct encodes the firefly luciferase reporter gene under the control of a minimal (m)CMV promoter and tandem repeats of the PPAR transcriptional response element. The number of response elements and the intervening sequence between these response elements has been experimentally optimized to maximize the signal to noise ratio. This construct monitors both increases and decreases in the transcriptional activity of PPAR. The constitutively expressing Renilla construct encodes the Renilla luciferase reporter gene under the control of a CMV immediately early enhancer/promoter and acts as an internal control for normalizing transfection efficiencies and monitoring cell viability. Using a simple dual-luciferase assay, you can easily monitor the activity of PPAR signaling pathways and determine the effect of various treatments, such as gene knockdown, over-expression, and chemical compounds on those pathways.

For further details, consult the *Cignal Reporter Assay Handbook*.

Shipping and storage

The Cignal Reporter Assay Kit is shipped on dry ice or blue ice. Store at -20°C. When stored under these conditions, this product can be used for 6 months from the date of receipt without reduction in performance.

Note: Open the package and store the products appropriately immediately on receipt.

Performance data

Visit the Website to view performance data for this product.



Contents

Product	Contents	Cat. no.
Signal Reporter Assay Kit	For 500 reporter assays, 500 negative control assays, and 250 positive control assays when using the recommended 96-well plate transfection protocol; 500 µl (100 ng/µl) Reporter (40:1 mixture of inducible firefly luciferase construct and constitutively expressing Renilla luciferase construct); 500 µl (100 ng/µl) Negative control (40:1 mixture of non-inducible firefly luciferase construct and constitutively expressing Renilla luciferase construct); 250 µl (100 ng/µl) Positive control (40:1:1 mixture of constitutively expressing GFP, constitutively expressing firefly luciferase, and constitutively expressing Renilla luciferase constructs)	Varies

Signal Reporter Assay Kit products are intended for molecular biology applications. These products are not intended for the diagnosis, prevention, or treatment of a disease.

For up-to-date licensing information and product-specific disclaimers, see the respective QIAGEN kit handbook or user manual. QIAGEN kit handbooks and user manuals are available at www.qiagen.com or can be requested from QIAGEN Technical Services or your local distributor.

Trademarks: QIAGEN® (QIAGEN Group).
1065997 03/2011 © 2011 QIAGEN, all rights reserved.

www.qiagen.com	Canada ■ 800-572-9613 China ■ 8621-3865-3865 Denmark ■ 80-885945 Australia ■ 1-800-243-800 Austria ■ 0800/281010 Belgium ■ 0800-79612 Brazil ■ 0800-557779	Ireland ■ 1800 555 049 Italy ■ 800-787980 Japan ■ 03-6890-7300 Finland ■ 0800-914416 France ■ 01-60-920-930 Germany ■ 02103-29-12000 Hong Kong ■ 800 933 965	Norway ■ 800-18859 Singapore ■ 1800-742-4368 Spain ■ 91-630-7050 Korea (South) ■ 080-000-7145 Luxembourg ■ 8002 2076 Mexico ■ 01-800-7742-436 The Netherlands ■ 0800 0229592	Norway ■ 800-18859 Singapore ■ 1800-742-4368 Spain ■ 91-630-7050 Sweden ■ 020-790282 Switzerland ■ 055-254-22-11 UK ■ 01293-422-911 USA ■ 800-426-8157
--	---	---	---	---



Sample & Assay Technologies