# RT<sup>2</sup> Profiler PCR Array (Rotor-Gene® Format) Human Hypoxia Signaling Pathway Plus

#### Cat. no. 330231 PAHS-032YR

#### For pathway expression analysis

Format	For use with the following real-time cyclers		
RT <sup>2</sup> Profiler PCR Array,	Rotor-Gene Q, other Rotor-Gene cyclers		
Format R			

#### Description

The Human Hypoxia Signaling Pathway Plus RT<sup>2</sup> Profiler PCR Array profiles the expression of 84 genes that respond to low oxygen levels. It also determines whether hypoxia pathway activity is increased or unchanged in experimental samples. Oxygen is required for aerobic energy metabolism processes such as oxidative phosphorylation. Low oxygen conditions activate the hypoxia signaling pathway in eukaryotic cells, primarily via the hypoxia inducible factor (HIF) transcription factor. HIF heterodimers consist of a constitutively-expressed beta subunit and one of 3 alpha subunit isoforms whose expression is tightly regulated. The presence of oxygen activates prolyl hydroxylases to hydroxylate HIF, leading to its polyubiquitination and degradation. Under low oxygen conditions, prolyl hydroxylase inactivity allows HIF to accumulate, initiating target gene expression. Hypoxia-inducible target genes mediate multiple biological functions, such as angiogenesis, hematopoiesis, and the maintenance of vascular tone to provide or replenish tissues with blood and oxygen. Hypoxia signaling dysregulation commonly occurs in diseases such as tumor angiogenesis and chronic inflammation. Hundreds of HIF target genes have been identified using experimental techniques such as expression studies and chromatin immunoprecipitation (ChIP) as well as bioinformatic analysis of predicted transcription factor binding sites. This array includes HIF signaling transcription factors, HIF interacting proteins, and highly relevant target genes identified by multiple studies. Results obtained with this array can be used to analyze activation or inhibition of hypoxia signaling. The array also includes 16 experimentally derived Signature Biomarker Genes which, along with classification algorithms, are used to generate the activity score. A set of controls present on each array enables data analysis using the ??CT method of relative quantification, assessment of reverse transcription performance, genomic DNA contamination, and PCR performance. Using real-time PCR, research studies can easily and reliably determine hypoxia signaling pathway activity and analyze the expression of a focused panel of genes related to the hypoxia signaling pathway with this array.

For further details, consult the RT<sup>2</sup> Profiler PCR Array Handbook.

## Shipping and storage

RT<sup>2</sup> Profiler PCR Arrays in the Rotor-Gene format are shipped at ambient temperature, on dry ice, or blue ice packs depending on destination and accompanying products.

For long term storage, keep plates at –20°C.

**Note:** Ensure that you have the correct RT<sup>2</sup> Profiler PCR Array format for your real-time cycler (see table above).

Note: Open the package and store the products appropriately immediately on receipt.



Sample & Assay Technologies

## Array layout

The 96 real-time assays in the Rotor-Gene format are located in wells 1–96 of the Rotor-Disc<sup>™</sup> (plate A1–A12=Rotor-Disc 1–12, plate B1–B12=Rotor-Disc 13–24, etc.). To maintain data analysis compatibility, wells 97–100 do not contain real-time assays but will contain master mix to account for weight balance.

## Gene table: RT<sup>2</sup> Profiler PCR Array

Position	UniGene	GenBank	Symbol	Description
A01	Hs.441047	NM 001124	ADM	Adrenomedullin
A02	Hs.167046	NM 000676	ADORA2B	Adenosine A2b receptor
A03	Hs.732822	NM 000034	ALDOA	Aldolase A, fructose-bisphosphate
A04	Hs.511605	NM 004039	ANXA2	Annexin A2
A05	Hs.73722	 NM 080649	APEX1	APEX nuclease (multifunctional DNA repair enzyme) 1
A06	Hs.632446	NM 001668	ARNT	Aryl hydrocarbon receptor nuclear translocator
A07	Hs.271791	NM 001184	ATR	Ataxia telangiectasia and Rad3 related
A08	Hs.725208	NM 000057	BLM	Bloom syndrome, RecQ helicase-like
A08 A09	Hs.740456	NM 004354	CCNG2	Cvclin G2
A09 A10	Hs.491912	NM 006837	COPS5	COP9 constitutive photomorphogenic homolog subunit 5 (Arabidopsis)
		NM 000308		
A11	Hs.609336	-	CTSA	Cathepsin A
A12	Hs.744875	NM_019058	DDIT4	DNA-damage-inducible transcript 4
B01	Hs.164419	NM_025219	DNAJC5	DnaJ (Hsp40) homolog, subfamily C, member 5
B02	Hs.713645	NM_001955	EDN1	Endothelin 1
B03	Hs.444450	NM_022051	EGLN1	Egl nine homolog 1 (C. elegans)
B04	Hs.730737	NM_053046	EGLN2	Egl nine homolog 2 (C. elegans)
B05	Hs.708393	NM_001964	EGR1	Early growth response 1
B06	Hs.411641	NM_004095	EIF4EBP1	Eukaryotic translation initiation factor 4E binding protein 1
B07	Hs.517145	NM_001428	ENO1	Enolase 1, (alpha)
B08	Hs.2303	NM_000799	EPO	Erythropoietin
B09	Hs.592304	NM 014584	ERO1L	ERO1-like (S. cerevisiae)
B10	Hs.361463	NM 000504	F10	Coagulation factor X
B11	Hs.62192	NM 001993	F3	Coagulation factor III (thromboplastin, tissue factor)
B12	Hs.25647	NM 005252	FOS	FBJ murine osteosarcoma viral oncogene homolog
C01	Hs.436062	NM 000158	GBE1	Glucan (1,4-alpha-), branching enzyme 1
C02	Hs.466471	NM 000175	GPI	Glucose-6-phosphate isomerase
C03	Hs.386225	NM 002103	GYS1	Glycogen synthase 1 (muscle)
000	113.300223	14/1_002103	0151	Hypoxia inducible factor 1, alpha subunit (basic helix-loop-helix transcription
C04	Hs.719495	NM_001530	HIF1A	factor)
C05	Hs.500788	NM 017902	HIF1AN	Hypoxia inducible factor 1, alpha subunit inhibitor
C06	Hs.420830	NM 152794	HIF3A	Hypoxia inducible factor 3, alpha subunit
C07	Hs.517581	NM 002133	HMOX1	Heme oxygenase (decycling) 1
C08	Hs.116462	 NM 178849	HNF4A	Hepatocyte nuclear factor 4, alpha
C09	Hs.76095	NM 003897	IER3	Immediate early response 3
C10	Hs.450230	NM 000598	IGFBP3	Insulin-like growth factor binding protein 3
C11	Hs.514505	NM 015167	JMJD6	Jumonji domain containing 6
C12	Hs.531081	NM 002306	LGALS3	Lectin, galactoside-binding, soluble, 3
D01	Hs.102267	NM 002317	LOALSS	
		-		Lysyl oxidase
D02	Hs.653654	NM_005921	MAP3K1	Mitogen-activated protein kinase kinase 1
D03	Hs.132966	NM_000245	MET	Met proto-oncogene (hepatocyte growth factor receptor)
D04	Hs.407995	NM_002415	MIF	Macrophage migration inhibitory factor (glycosylation-inhibiting factor)
D05	Hs.297413	NM_004994	MMP9	Matrix metallopeptidase 9 (gelatinase B, 92kDa gelatinase, 92kDa type IV collagenase)
D06	Hs.728542	NM 005962	MXI1	MAX interactor 1
D07	Hs.489615	NM 005746	NAMPT	Nicotinamide phosphoribosyltransferase
D08	Hs.596314	NM 003743	NCOA1	Nuclear receptor coactivator 1
D00	Hs.618430	NM 003998	NFKB1	Nuclear factor of kappa light polypeptide gene enhancer in B-cells 1
D07	Hs.647092	NM 000603	NOS3	Nitric oxide synthase 3 (endothelial cell)
D10	Hs.467701	NM 002539	ODC1	Ornithine decarboxylase 1
D11 D12	Hs.467701 Hs.464336	NM 000918	P4HB	,
				Prolyl 4-hydroxylase, beta polypeptide
E01	Hs.733780	NM_002610	PDK1	Pyruvate dehydrogenase kinase, isozyme 1
E02	Hs.445534	NM_002616	PER1	Period homolog 1 (Drosophila)
E03	Hs.195471	NM_004566	PFKFB3	6-phosphofructo-2-kinase/fructose-2,6-biphosphatase 3
E04	Hs.255093	NM_002626	PFKL	Phosphofructokinase, liver
E05	Hs.26010	NM_002627	PFKP	Phosphofructokinase, platelet
E06	Hs.252820	NM_002632	PGF	Placental growth factor
E07	Hs.81170	NM 002648	PIM1	Pim-1 oncogene

Position	UniGene	GenBank	Symbol	Description	
E08	Hs.534770	NM_002654	PKM	Pyruvate kinase, muscle	
E09	Hs.77274	NM_002658	PLAU	Plasminogen activator, urokinase	
E10	Hs.479396	NM_005349	RBPJ	Recombination signal binding protein for immunoglobulin kappa J region	
E11	Hs.515846	NM_006666	RUVBL2	RuvB-like 2 (E. coli)	
E12	Hs.713079	NM_000602	SERPINE1	Serpin peptidase inhibitor, clade E (nexin, plasminogen activator inhibitor type 1), member 1	
F01	Hs.500761	NM_004207	SLC16A3	Solute carrier family 16, member 3 (monocarboxylic acid transporter 4)	
F02	Hs.473721	NM_006516	SLC2A1	Solute carrier family 2 (facilitated glucose transporter), member 1	
F03	Hs.529618	NM_003234	TFRC	Transferrin receptor (p90, CD71)	
F04	Hs.740601	NM_000546	TP53	Tumor protein p53	
F05	Hs.524219	NM_000365	TPI1	Triosephosphate isomerase 1	
F06	Hs.709057	NM_006472	TXNIP	Thioredoxin interacting protein	
F07	Hs.454534	NM 003367	USF2	Upstream transcription factor 2, c-fos interacting	
F08	Hs.519320	NM_003374	VDAC1	Voltage-dependent anion channel 1	
F09	Hs.155247	NM 005165	ALDOC	Aldolase C, fructose-bisphosphate	
F10	Hs.9613	NM 001039667	ANGPTL4	Angiopoietin-like 4	
F11	Hs.508154	NM 181726	ANKRD37	Ankyrin repeat domain 37	
F12	Hs.744856	NM 003670	BHLHE40	Basic helix-loop-helix family, member e40	
G01	Hs.144873	NM 004052	BNIP3	BCL2/adenovirus E1B 19kDa interacting protein 3	
G02	Hs.131226	NM 004331	<b>BNIP3L</b>	BCL2/adenovirus E1B 19kDa interacting protein 3-like	
G03	Hs.63287	NM 001216	CA9	Carbonic anhydrase IX	
G04	Hs.584881	NM 014367	FAM162A	Family with sequence similarity 162, member A	
G05	Hs.591588	NM_000189	HK2	Hexokinase 2	
G06	Hs.2795	NM 005566	LDHA	Lactate dehydrogenase A	
G07	Hs.618002	NM 006096	NDRG1	N-myc downstream regulated 1	
G08	Hs.593005	NM 000917	P4HA1	Prolyl 4-hydroxylase, alpha polypeptide l	
G09	Hs.476217	NM 004567	PFKFB4	6-phosphofructo-2-kinase/fructose-2,6-biphosphatase 4	
G10	Hs.78771	NM 000291	PGK1	Phosphoglycerate kinase 1	
G11	Hs.419240	NM 006931	SLC2A3	Solute carrier family 2 (facilitated glucose transporter), member 3	
G12	Hs.73793	NM 003376	VEGFA	Vascular endothelial growth factor A	
H01	Hs.520640	NM 001101	ACTB	Actin, beta	
H02	Hs.534255	NM 004048	B2M	Beta-2-microglobulin	
H03	Hs.544577	NM 002046	GAPDH	Glyceraldehyde-3-phosphate dehydrogenase	
H04	Hs.412707	NM 000194	HPRT1	Hypoxanthine phosphoribosyltransferase 1	
H05	Hs.546285	NM 001002	RPLPO	Ribosomal protein, large, P0	
H06	N/A	SA 00105	HGDC	Human Genomic DNA Contamination	
H07	N/A	SA 00104	RTC	Reverse Transcription Control	
H08	N/A	SA 00104	RTC	Reverse Transcription Control	
H09	N/A	SA 00104	RTC	Reverse Transcription Control	
H10	N/A	SA 00103	PPC	Positive PCR Control	
H11	N/A	SA 00103	PPC	Positive PCR Control	
H12	N/A	SA 00103	PPC	Positive PCR Control	

## **Related products**

For optimal performance, RT<sup>2</sup> Profiler PCR Arrays should be used together with the RT<sup>2</sup> First Strand Kit for cDNA synthesis and RT<sup>2</sup> SYBR<sup>®</sup> Green qPCR Mastermixes for PCR.

Product	Contents	Cat. no.
RT <sup>2</sup> First Strand Kit (12)	Enzymes and reagents for cDNA synthesis	330401
RT² SYBR Green ROX™ FAST Mastermix (2)*	For 2 x 96 assays in 96-well plates; suitable for use with the Rotor-Gene Q and other Rotor-Gene cyclers	330620

\* Larger kit sizes available; please inquire.

RT<sup>2</sup> Profiler PCR Array products are intended for molecular biology applications. These products are not intended for the diagnosis, prevention, or treatment of a disease.

For up-to-date licensing information and product-specific disclaimers, see the respective QIAGEN kit handbook or user manual. QIAGEN kit handbooks and user manuals are available at <u>www.qiagen.</u> <u>com</u> or can be requested from QIAGEN Technical Services or your local distributor.

Trademarks: QIAGEN<sup>®</sup>, Rotor-Gene<sup>®</sup>, Rotor-Disc<sup>™</sup> (QIAGEN Group); ROX<sup>™</sup> (Applera Corporation or its subsidiaries); SYBR<sup>®</sup> (Molecular Probes, Inc.).

1067688 03/2011 © 2011 QIAGEN, all rights reserved.

www.qiagen.com	<b>Canada =</b> 800-572-9613	Ireland • 1800 555 049	Norway = 800-18859	
	China • 8621-3865-3865	Italy = 800-787980	Singapore = 1800-742-4368	
	<b>Denmark</b> • 80-885945	Japan = 03-6890-7300	<b>Spain</b> ■ 91-630-7050	$\bullet \bullet \bullet \bullet \bullet \bullet \bullet$
Australia • 1-800-243-800	Finland • 0800-914416	Korea (South) = 080-000-7145	Sweden = 020-790282	
Austria = 0800/281010	France = 01-60-920-930	Luxembourg = 8002 2076	Switzerland • 055-254-22-11	
Belgium = 0800-79612	Germany • 02103-29-12000	<b>Mexico</b> • 01-800-7742-436	UK • 01293-422-911	QIAGEN
Brazil • 0800-557779	Hong Kong  ■ 800 933 965	<b>The Netherlands</b> • 0800 0229592	<b>USA •</b> 800-426-8157	GIAGEN

Sample & Assay Technologies