RT² Profiler PCR Array (Rotor-Gene® Format) Human Retinoic Acid Signaling

Cat. no. 330231 PAHS-180ZR

For pathway expression analysis

| Format | For use with the following real-time cyclers |
|-------------------------------------|--|
| RT ² Profiler PCR Array, | Rotor-Gene Q, other Rotor-Gene cyclers |
| Format R | |

Description

The Human Retinoic Acid Pathway RT² Profiler PCR Array profiles the expression of 84 key genes involved in retinoic acid signaling. Retinoic acid (RA) is the primary functional derivative of vitamin A (retinol) and its activity is implicated in many aspects of vertebrate development and homeostasis, while disruptions in this pathway cause developmental abnormalities and disrupt function in adipose, cardiac, nervous, reproductive, and integumentary tissues, among others. RA acts primarily by binding a family of nuclear receptors (the retinoic acid receptors alpha, beta, and gamma) that then heterodimerize with their partners (the retinoid X receptors alpha, beta, and gamma) and alter transcription. Additionally, recent evidence suggests that an alternative receptor (PPAR delta) also responds to RA signaling in some tissues and that retinol, and RA may also induce non-genomic cellular effects. Thus, the effect of RA depends on the expression of its cognate receptors, enzymes responsible for converting retinol to RA, metabolic enzymes that reduce RA levels, and transport proteins, some of which act as signal transducers that appear to influence the distribution of RA and its metabolic precursors. This array includes genes encoding the known receptors and proteins responsible for synthesis, transport and metabolism of RA and its precursors, as well as downstream targets of RA signaling. Using real-time PCR, research studies can easily and reliably analyze the expression of a focused panel of genes involved in retinoic acid signaling with this array.

For further details, consult the RT² Profiler PCR Array Handbook.

Shipping and storage

RT² Profiler PCR Arrays in the Rotor-Gene format are shipped at ambient temperature, on dry ice, or blue ice packs depending on destination and accompanying products.

For long term storage, keep plates at -20°C.

Note: Ensure that you have the correct RT² Profiler PCR Array format for your real-time cycler (see table above).

Note: Open the package and store the products appropriately immediately on receipt.



Array layout

The 96 real-time assays in the Rotor-Gene format are located in wells 1–96 of the Rotor-Disc[™] (plate A1–A12=Rotor-Disc 1–12, plate B1–B12=Rotor-Disc 13–24, etc.). To maintain data analysis compatibility, wells 97–100 do not contain real-time assays but will contain master mix to account for weight balance.

Gene table: RT² Profiler PCR Array

| Position | UniGene | GenBank | Symbol | Description | |
|------------|-----------|-----------|---------|---|--|
| A01 | Hs.654433 | NM_000667 | ADH1A | Alcohol dehydrogenase 1A (class I), alpha polypeptide | |
| A02 | Hs.76392 | NM_000689 | ALDH1A1 | Aldehyde dehydrogenase 1 family, member A1 | |
| A03 | Hs.643455 | NM_003888 | ALDH1A2 | Aldehyde dehydrogenase 1 family, member A2 | |
| A04 | Hs.459538 | NM_000693 | ALDH1A3 | Aldehyde dehydrogenase 1 family, member A3 | |
| A05 | Hs.237658 | NM_001643 | APOA2 | Apolipoprotein A-II | |
| A06 | Hs.703025 | NM_004316 | ASCL1 | Achaete-scute complex homolog 1 (Drosophila) | |
| A07 | Hs.744856 | NM 003670 | BHLHE40 | Basic helix-loop-helix family, member e40 | |
| A08 | Hs.73853 | NM 001200 | BMP2 | Bone morphogenetic protein 2 | |
| A09 | Hs.479214 | NM 001775 | CD38 | CD38 molecule | |
| A10 | Hs.1545 | NM 001804 | CDX1 | Caudal type homeobox 1 | |
| A11 | Hs.733236 | NM 017780 | CHD7 | Chromodomain helicase DNA binding protein 7 | |
| A12 | Hs.346950 | NM 004378 | CRABP1 | Cellular retinoic acid binding protein 1 | |
| B01 | Hs.405662 | NM 001878 | CRABP2 | Cellular retinoic acid binding protein 2 | |
| B02 | Hs.154654 | NM 000104 | CYP1B1 | Cytochrome P450, family 1, subfamily B, polypeptide 1 | |
| B03 | Hs.150595 | NM 000783 | CYP26A1 | Cytochrome P450, family 26, subfamily A, polypeptide 1 | |
| B04 | Hs.91546 | NM 019885 | CYP26B1 | Cytochrome P450, family 26, subfamily B, polypeptide 1 | |
| B05 | Hs.369993 | NM 183374 | CYP26C1 | Cytochrome P450, family 26, subfamily C, polypeptide 1 | |
| B06 | Hs.34780 | NM 178153 | DCX | Doublecortin | |
| B07 | Hs.289347 | NM 004753 | DHRS3 | Dehydrogenase/reductase (SDR family) member 3 | |
| B08 | Hs.179608 | NM 005771 | DHRS9 | Dehydrogenase/reductase (SDR family) member 9 | |
| B09 | Hs.99348 | NM 005221 | DLX5 | Distal-less homeobox 5 | |
| B10 | Hs.144700 | NM_004429 | EFNB1 | Ephrin-B1 | |
| B11 | Hs.708393 | NM 001964 | EGR1 | Early growth response 1 | |
| B12 | Hs.2303 | NM 000799 | EPO | Erythropoietin | |
| C01 | Hs.408061 | NM 001444 | FABP5 | Fatty acid binding protein 5 (psoriasis-associated) | |
| C02 | Hs.57710 | NM 006119 | FGF8 | Fibroblast growth factor 8 (androgen-induced) | |
| C03 | Hs.163484 | NM 004496 | FOXA1 | Forkhead box A1 | |
| C04 | Hs.741222 | NM 005249 | FOXG1 | Forkhead box G1 | |
| C05 | Hs.243987 | NM 002052 | GATA4 | GATA binding protein 4 | |
| C06 | Hs.184945 | NM 001485 | GBX2 | Gastrulation brain homeobox 2 | |
| C07 | Hs.632702 | NM 005269 | GLI1 | GLI family zinc finger 1 | |
| C08 | Hs.191144 | NM_000458 | HNF1B | HNF1 homeobox B | |
| C09 | Hs.67397 | NM 005522 | HOXA1 | Homeobox A1 | |
| C10 | Hs.655218 | NM 019102 | HOXA5 | Homeobox A5 | |
| C11 | Hs.99992 | NM 002144 | HOXB1 | Homeobox B1 | |
| C12 | Hs.664706 | NM 024015 | HOXB4 | Homeobox B4 | |
| D01 | Hs.162795 | NM 002153 | HSD17B2 | Hydroxysteroid (17-beta) dehydrogenase 2 | |
| D02 | Hs.505 | NM 002202 | ISL1 | ISL LIM homeobox 1 | |
| D03 | Hs.626544 | NM 000214 | JAG1 | Jagged 1 | |
| D04 | Hs.376206 | NM 004235 | KLF4 | Kruppel-like factor 4 (gut) | |
| D05 | Hs.656214 | NM 020997 | LEFTY1 | Left-right determination factor 1 | |
| D06 | Hs.443727 | NM 005568 | LHX1 | LIM homeobox 1 | |
| D07 | Hs.658427 | NM 004744 | LRAT | Lecithin retinol acyltransferase (phosphatidylcholineretinol O-acyltransferas | |
| D08 | Hs.169487 | NM 005461 | MAFB | V-maf musculoaponeurotic fibrosarcoma oncogene homolog B (avian) | |
| D09 | Hs.510989 | NM 172316 | MEIS2 | Meis homeobox 2 | |
| D10 | Hs.89404 | NM 002449 | MSX2 | Msh homeobox 2 | |
| D11 | Hs.202453 | NM 002467 | MYC | V-myc myelocytomatosis viral oncogene homolog (avian) | |
| D12 | Hs.635882 | NM 024865 | NANOG | Nanog homeobox | |
| E01 | Hs.709709 | NM 002500 | NEUROD1 | Neurogenic differentiation 1 | |
| E02 | Hs.155017 | NM 003489 | NRIP1 | Nuclear receptor interacting protein 1 | |
| E03 | Hs.732068 | NM 005806 | OLIG2 | Oligodendrocyte lineage transcription factor 2 | |
| | Hs.741558 | NM 021728 | OTX2 | Orthodenticle homeobox 2 | |
| FO4 | | NM 000280 | PAX6 | Paired box 6 | |
| E04 | He 611274 | | | | |
| E05 | Hs.611376 | | | | |
| E05 E06 | Hs.738484 | NM_000325 | PITX2 | Paired-like homeodomain 2 | |
| E05 | | | | | |

| Position | UniGene | GenBank | Symbol | Description | |
|--------------------------|-------------------|----------------------------------|--------------------------|--|--|
| E10 | Hs.162646 | NM_015869 | PPARG | Peroxisome proliferator-activated receptor gamma | |
| E11 | Hs.654583 | NM_000964 | RARA | Retinoic acid receptor, alpha | |
| E12 | Hs.733004 | NM_000965 | RARB | Retinoic acid receptor, beta | |
| F01 | Hs.733399 | NM_000966 | RARG | Retinoic acid receptor, gamma | |
| F02 | Hs.17466 | NM_004585 | RARRES3 | Retinoic acid receptor responder (tazarotene induced) 3 | |
| F03 | Hs.529571 | NM_002899 | RBP1 | Retinol binding protein 1, cellular | |
| F04 | Hs.655516 | NM_004164 | RBP2 | Retinol binding protein 2, cellular | |
| F05 | Hs.50223 | NM_006744 | RBP4 | Retinol binding protein 4, plasma | |
| F06 | Hs.244940 | NM_172037 | RDH10 | Retinol dehydrogenase 10 (all-trans) | |
| F07 | Hs.350321 | NM_020630 | RET | Ret proto-oncogene | |
| F08 | Hs.590886 | NM_002957 | RXRA | Retinoid X receptor, alpha | |
| F09 | Hs.388034 | NM 021976 | RXRB | Retinoid X receptor, beta | |
| F10 | Hs.26550 | NM 006917 | RXRG | Retinoid X receptor, gamma | |
| F11 | Hs.164537 | NM 000193 | SHH | Sonic hedgehog | |
| F12 | Hs.732963 | NM 003106 | SOX2 | SRY (sex determining region Y)-box 2 | |
| G01 | Hs.647409 | NM 000346 | SOX9 | SRY (sex determining region Y)-box 9 | |
| G02 | Hs.733635 | NM 004176 | SREBF1 | Sterol regulatory element binding transcription factor 1 | |
| G03 | Hs.24553 | NM 022369 | STRA6 | Stimulated by retinoic acid gene 6 homolog (mouse) | |
| G04 | Hs.592279 | NM 182489 | STRA8 | Stimulated by retinoic acid gene 8 homolog (mouse) | |
| G05 | Hs.173984 | NM 005992 | TBX1 | T-box 1 | |
| G06 | Hs.473152 | NM_003222 | TFAP2C | Transcription factor AP-2 gamma (activating enhancer binding protein 2 | |
| G07 | Hs.133379 | NM 003238 | TGFB2 | Transforming growth factor, beta 2 | |
| | | NM_004613 | TGM2 | Transglutaminase 2 (C polypeptide, | |
| G08 | Hs.517033 | | | protein-glutamine-gamma-glutamyltransferase) | |
| G09 | Hs.511743 | NM 006086 | TUBB3 | Tubulin, beta 3 | |
| G10 | Hs.249211 | NM 021833 | UCP1 | Uncoupling protein 1 (mitochondrial, proton carrier) | |
| G11 | Hs.643085 | NM 003392 | WNT5A | Wingless-type MMTV integration site family, member 5A | |
| G12 | Hs.591274 | NM 058244 | WNT8A | Wingless-type MMTV integration site family, member 8A | |
| H01 | Hs.520640 | NM 001101 | ACTB | Actin, beta | |
| H02 | Hs.534255 | NM 004048 | B2M | Beta-2-microalobulin | |
| H03 | Hs.544577 | NM 002046 | GAPDH | Glyceraldehyde-3-phosphate dehydrogenase | |
| H04 | Hs.412707 | NM 000194 | HPRT1 | Hypoxanthine phosphoribosyltransferase 1 | |
| H05 | Hs.546285 | NM 001002 | RPLPO | Ribosomal protein, large, PO | |
| H06 | N/A | SA 00105 | HGDC | Human Genomic DNA Contamination | |
| H07 | N/A | SA 00104 | RTC | Reverse Transcription Control | |
| H08 | N/A | SA 00104 | RTC | · | |
| H09 | N/A | SA 00104 | RTC | · · | |
| H10 | N/A | SA 00103 | PPC | ' | |
| H11 | N/A | SA 00103 | PPC | | |
| H12 | | _ | PPC | | |
| H08 H09 H10 H11 | N/A N/A N/A | SA_00104 SA_00104 SA_00103 | RTC RTC PPC PPC | Reverse Transcription Control Reverse Transcription Control Reverse Transcription Control Positive PCR Control Positive PCR Control Positive PCR Control | |

Related products

For optimal performance, RT² Profiler PCR Arrays should be used together with the RT² First Strand Kit for cDNA synthesis and RT² SYBR[®] Green qPCR Mastermixes for PCR.

| Product | Contents | Cat. no. |
|--|--|----------|
| RT ² First Strand Kit (12) | Enzymes and reagents for cDNA synthesis | 330401 |
| RT ² SYBR Green ROX [™] FAST Mastermix (2)* | For 2 x 96 assays in 96-well plates; suitable for use with the Rotor-Gene Q and other Rotor-Gene cyclers | 330620 |

^{*} Larger kit sizes available; please inquire.

RT² Profiler PCR Array products are intended for molecular biology applications. These products are not intended for the diagnosis, prevention, or treatment of a disease.

For up-to-date licensing information and product-specific disclaimers, see the respective QIAGEN kit handbook or user manual. QIAGEN kit handbooks and user manuals are available at www.qiagen. com or can be requested from QIAGEN Technical Services or your local distributor.

Trademarks: QIAGEN®, Rotor-Gene®, Rotor-Disc™ (QIAGEN Group); ROX™ (Applera Corporation or its subsidiaries); SYBR® (Molecular Probes, Inc.).

1067688 03/2011 © 2011 QIAGEN, all rights reserved.

www.aiaaen.com Australia • 1-800-243-800 Austria • 0800/281010 Belgium • 0800-79612 Brazil • 0800-557779

Canada • 800-572-9613 China • 8621-3865-3865 Denmark ■ 80-885945 Finland • 0800-914416 France • 01-60-920-930 Germany • 02103-29-12000 Hong Kong • 800 933 965

Ireland • 1800 555 049 Italy • 800-787980 Japan ■ 03-6890-7300 Korea (South) • 080-000-7145 Luxembourg ■ 8002 2076 Mexico • 01-800-7742-436 The Netherlands • 0800 0229592 USA • 800-426-8157

Norway • 800-18859 Singapore ■ 1800-742-4368 Spain ■ 91-630-7050 Sweden • 020-790282 Switzerland ■ 055-254-22-11 UK • 01293-422-911

