

RT² Profiler PCR Array (Rotor-Gene[®] Format)

Mouse Estrogen Receptor Signaling

Cat. no. 330231 PAMM-005ZR

For pathway expression analysis

| Format | For use with the following real-time cyclers |
|-------------------------------------|--|
| RT ² Profiler PCR Array, | Rotor-Gene Q, other Rotor-Gene cyclers |
| Format R | |

Description

The Mouse Estrogen Receptor Signaling RT² Profiler PCR Array profiles the expression of 84 key genes involved in estrogen receptor (ER) activation and response. Estrogen receptors are steroid hormone receptors important in development, growth, and reproduction. The 2 well-characterized ERs, alpha and beta, interact with a variety of co-regulators in the nucleus and initiate target gene transcription. ERs can also associate with the cell membrane, where they activate downstream signaling pathways. ERs play a large role in female organ cancers, especially breast cancer. The mechanisms of ER signaling are not entirely understood since tissue-specific ER responses vary depending on the agonist or antagonist. ER signaling is important in mammalian systems other than female organs. For example, ERs are involved in neurological development, and estrogens play a protective role against cardiovascular disease and osteoporosis, although the exact mechanisms of these processes are under investigation. This array includes the ERs, their co-regulators and interacting proteins, and downstream target genes. Results obtained with this array can yield insights into ER mechanisms and responses. Using real-time PCR, research studies can easily and reliably analyze the expression of a focused panel of genes involved in estrogenic signaling with this array.

For further details, consult the *RT² Profiler PCR Array Handbook*.

Shipping and storage

RT² Profiler PCR Arrays in the Rotor-Gene format are shipped at ambient temperature, on dry ice, or blue ice packs depending on destination and accompanying products.

For long term storage, keep plates at -20°C.

Note: Ensure that you have the correct RT² Profiler PCR Array format for your real-time cycler (see table above).

Note: Open the package and store the products appropriately immediately on receipt.



Sample & Assay Technologies

Array layout

The 96 real-time assays in the Rotor-Gene format are located in wells 1–96 of the Rotor-Disc™ (plate A1–A12=Rotor-Disc 1–12, plate B1–B12=Rotor-Disc 13–24, etc.). To maintain data analysis compatibility, wells 97–100 do not contain real-time assays but will contain master mix to account for weight balance.

Gene table: RT² Profiler PCR Array

| Position | UniGene | GenBank | Symbol | Description |
|----------|-----------|--------------|----------|--|
| A01 | Mm.298908 | NM_001008533 | Adora1 | Adenosine A1 receptor |
| A02 | Mm.341377 | NM_013464 | Ahr | Aryl-hydrocarbon receptor |
| A03 | Mm.2969 | NM_009648 | Akap1 | A kinase (PRKA) anchor protein 1 |
| A04 | Mm.38469 | NM_009685 | Apbb1 | Amyloid beta (A4) precursor protein-binding, family B, member 1 |
| A05 | Mm.3758 | NM_009954 | Bcar1 | Breast cancer anti-estrogen resistance 1 |
| A06 | Mm.238213 | NM_009743 | Bcl2l1 | Bcl2-like 1 |
| A07 | Mm.1442 | NM_007540 | Bdnf | Brain derived neurotrophic factor |
| A08 | Mm.6813 | NM_007554 | Bmp4 | Bone morphogenetic protein 4 |
| A09 | Mm.595 | NM_007557 | Bmp7 | Bone morphogenetic protein 7 |
| A10 | Mm.244975 | NM_009764 | Brcal | Breast cancer 1 |
| A11 | Mm.19131 | NM_009778 | C3 | Complement component 3 |
| A12 | Mm.28278 | NM_007616 | Cav1 | Caveolin 1, caveolae protein |
| B01 | Mm.867 | NM_011331 | Ccl12 | Chemokine (C-C motif) ligand 12 |
| B02 | Mm.273049 | NM_007631 | Ccnd1 | Cyclin D1 |
| B03 | Mm.272321 | NM_010828 | Cited2 | Cbp/p300-interacting transactivator, with Glu/Asp-rich carboxy-terminal domain, 2 |
| B04 | Mm.16831 | NM_021273 | Ckb | Creatine kinase, brain |
| B05 | Mm.390287 | NM_010217 | Ctgf | Connective tissue growth factor |
| B06 | Mm.231395 | NM_009983 | Ctsd | Cathepsin D |
| B07 | Mm.5199 | NM_007810 | Cyp19a1 | Cytochrome P450, family 19, subfamily a, polypeptide 1 |
| B08 | Mm.14089 | NM_009992 | Cyp1a1 | Cytochrome P450, family 1, subfamily a, polypeptide 1 |
| B09 | Mm.477528 | NM_019480 | Ebag9 | Estrogen receptor-binding fragment-associated gene 9 |
| B10 | Mm.401670 | NM_010109 | Efna5 | Ephrin A5 |
| B11 | Mm.103737 | NM_018781 | Egr3 | Early growth response 3 |
| B12 | Mm.290822 | NM_001003817 | Erbb2 | V-erb-b2 erythroblastic leukemia viral oncogene homolog 2, neuro/glioblastoma derived oncogene homolog (avian) |
| C01 | Mm.373043 | NM_010153 | Erbb3 | V-erb-b2 erythroblastic leukemia viral oncogene homolog 3 (avian) |
| C02 | Mm.9213 | NM_007956 | Esr1 | Estrogen receptor 1 (alpha) |
| C03 | Mm.2561 | NM_010157 | Esr2 | Estrogen receptor 2 (beta) |
| C04 | Mm.246513 | NM_010234 | Fos | FBJ osteosarcoma oncogene |
| C05 | Mm.4578 | NM_008259 | Foxa1 | Forkhead box A1 |
| C06 | Mm.4913 | NM_008046 | Fst | Follistatin |
| C07 | Mm.27210 | NM_008062 | G6pdx | Glucose-6-phosphate dehydrogenase X-linked |
| C08 | Mm.1843 | NM_010480 | Hsp90aa1 | Heat shock protein 90, alpha (cytosolic), class A member 1 |
| C09 | Mm.268521 | NM_010512 | Igf1 | Insulin-like growth factor 1 |
| C10 | Mm.233799 | NM_010517 | Igfbp4 | Insulin-like growth factor binding protein 4 |
| C11 | Mm.405761 | NM_010518 | Igfbp5 | Insulin-like growth factor binding protein 5 |
| C12 | Mm.4952 | NM_010570 | Irs1 | Insulin receptor substrate 1 |
| D01 | Mm.1167 | NM_008416 | Junb | Jun-B oncogene |
| D02 | Mm.482691 | NM_008455 | Klkb1 | Kallikrein B, plasma 1 |
| D03 | Mm.260568 | NM_008478 | L1cam | L1 cell adhesion molecule |
| D04 | Mm.43831 | NM_008495 | Lgals1 | Lectin, galactose binding, soluble 1 |
| D05 | Mm.1514 | NM_008509 | Lpl | Lipoprotein lipase |
| D06 | Mm.269747 | NM_019919 | Libp1 | Latent transforming growth factor beta binding protein 1 |
| D07 | Mm.86646 | NM_010755 | Maff | V-maf musculoaponeurotic fibrosarcoma oncogene family, protein F (avian) |
| D08 | Mm.12926 | NM_013634 | Med1 | Mediator complex subunit 1 |
| D09 | Mm.4406 | NM_013599 | Mmp9 | Matrix metallopeptidase 9 |
| D10 | Mm.212577 | NM_054081 | Mta1 | Metastasis associated 1 |
| D11 | Mm.2444 | NM_010849 | Myc | Myelocytomatosis oncogene |
| D12 | Mm.336898 | NM_008668 | Nab2 | Ngfi-A binding protein 2 |
| E01 | Mm.301039 | NM_010881 | Ncoa1 | Nuclear receptor coactivator 1 |
| E02 | Mm.2537 | NM_008678 | Ncoa2 | Nuclear receptor coactivator 2 |
| E03 | Mm.476883 | NM_008679 | Ncoa3 | Nuclear receptor coactivator 3 |
| E04 | Mm.271814 | NM_011308 | Ncor1 | Nuclear receptor co-repressor 1 |
| E05 | Mm.278646 | NM_011424 | Ncor2 | Nuclear receptor co-repressor 2 |
| E06 | Mm.5167 | NM_010930 | Nov | Nephroblastoma overexpressed gene |
| E07 | Mm.5180 | NM_007430 | Nr0b1 | Nuclear receptor subfamily 0, group B, member 1 |

| Position | UniGene | GenBank | Symbol | Description |
|----------|-----------|--------------|----------|--|
| E08 | Mm.346759 | NM_011850 | Nr0b2 | Nuclear receptor subfamily 0, group B, member 2 |
| E09 | Mm.28989 | NM_010150 | Nr2f6 | Nuclear receptor subfamily 2, group F, member 6 |
| E10 | Mm.129481 | NM_008173 | Nr3c1 | Nuclear receptor subfamily 3, group C, member 1 |
| E11 | Mm.16794 | NM_030676 | Nr5a2 | Nuclear receptor subfamily 5, group A, member 2 |
| E12 | Mm.74711 | NM_173440 | Nrip1 | Nuclear receptor interacting protein 1 |
| F01 | Mm.271745 | NM_008737 | Nrp1 | Neuropilin 1 |
| F02 | Mm.482226 | NM_021517 | Pdzk1 | PDZ domain containing 1 |
| F03 | Mm.340601 | NM_029231 | Pelp1 | Proline, glutamic acid and leucine rich protein 1 |
| F04 | Mm.12798 | NM_008829 | Pgr | Progesterone receptor |
| F05 | Mm.36241 | NM_007531 | Phb2 | Prohibitin 2 |
| F06 | Mm.228798 | NM_008957 | Ptch1 | Patched homolog 1 |
| F07 | Mm.292547 | NM_011198 | Ptg52 | Prostaglandin-endoperoxide synthase 2 |
| F08 | Mm.27348 | NM_019491 | Rala | V-rat simian leukemia viral oncogene homolog A (ras related) |
| F09 | Mm.439744 | NM_009024 | Rara | Retinoic acid receptor, alpha |
| F10 | Mm.100144 | NM_011313 | S100a6 | S100 calcium binding protein A6 (calcyclin) |
| F11 | Mm.255066 | NM_001163300 | Saffb | Scaffold attachment factor B |
| F12 | Mm.2093 | NM_011427 | Snai1 | Snail homolog 1 (Drosophila) |
| G01 | Mm.3468 | NM_007707 | Socs3 | Suppressor of cytokine signaling 3 |
| G02 | Mm.288474 | NM_009263 | Spp1 | Secreted phosphoprotein 1 |
| G03 | Mm.2854 | NM_009362 | Tff1 | Trefoil factor 1 |
| G04 | Mm.137222 | NM_031199 | Tgfa | Transforming growth factor alpha |
| G05 | Mm.3992 | NM_009368 | Tgfb3 | Transforming growth factor, beta 3 |
| G06 | Mm.4159 | NM_011580 | Thbs1 | Thrombospondin 1 |
| G07 | Mm.245084 | NM_009504 | Vdr | Vitamin D receptor |
| G08 | Mm.282184 | NM_009505 | Vegfa | Vascular endothelial growth factor A |
| G09 | Mm.13828 | NM_016873 | Wisp2 | WNT1 inducible signaling pathway protein 2 |
| G10 | Mm.20355 | NM_009523 | Wnt4 | Wingless-related MMTV integration site 4 |
| G11 | Mm.287544 | NM_009524 | Wnt5a | Wingless-related MMTV integration site 5A |
| G12 | Mm.469937 | NM_013842 | Xbp1 | X-box binding protein 1 |
| H01 | Mm.328431 | NM_007393 | Actb | Actin, beta |
| H02 | Mm.163 | NM_009735 | B2m | Beta-2 microglobulin |
| H03 | Mm.343110 | NM_008084 | Gapdh | Glyceraldehyde-3-phosphate dehydrogenase |
| H04 | Mm.3317 | NM_010368 | Gusb | Glucuronidase, beta |
| H05 | Mm.2180 | NM_008302 | Hsp90ab1 | Heat shock protein 90 alpha (cytosolic), class B member 1 |
| H06 | N/A | SA_00106 | MGDC | Mouse Genomic DNA Contamination |
| H07 | N/A | SA_00104 | RTC | Reverse Transcription Control |
| H08 | N/A | SA_00104 | RTC | Reverse Transcription Control |
| H09 | N/A | SA_00104 | RTC | Reverse Transcription Control |
| H10 | N/A | SA_00103 | PPC | Positive PCR Control |
| H11 | N/A | SA_00103 | PPC | Positive PCR Control |
| H12 | N/A | SA_00103 | PPC | Positive PCR Control |

Related products

For optimal performance, RT² Profiler PCR Arrays should be used together with the RT² First Strand Kit for cDNA synthesis and RT² SYBR[®] Green qPCR Mastermixes for PCR.

| Product | Contents | Cat. no. |
|---|--|----------|
| RT ² First Strand Kit (12) | Enzymes and reagents for cDNA synthesis | 330401 |
| RT ² SYBR Green ROX [™] FAST Mastermix (2)* | For 2 x 96 assays in 96-well plates; suitable for use with the Rotor-Gene Q and other Rotor-Gene cyclers | 330620 |

* Larger kit sizes available; please inquire.

RT² Profiler PCR Array products are intended for molecular biology applications. These products are not intended for the diagnosis, prevention, or treatment of a disease.

For up-to-date licensing information and product-specific disclaimers, see the respective QIAGEN kit handbook or user manual. QIAGEN kit handbooks and user manuals are available at www.qiagen.com or can be requested from QIAGEN Technical Services or your local distributor.

Trademarks: QIAGEN[®], Rotor-Gene[®], Rotor-Disc[™] (QIAGEN Group); ROX[™] (Applera Corporation or its subsidiaries); SYBR[®] (Molecular Probes, Inc.).

1067688 03/2011 © 2011 QIAGEN, all rights reserved.

www.qiagen.com

Australia ■ 1-800-243-800

Austria ■ 0800/281010

Belgium ■ 0800-79612

Brazil ■ 0800-557779

Canada ■ 800-572-9613

China ■ 8621-3865-3865

Denmark ■ 80-885945

Finland ■ 0800-914416

France ■ 01-60-920-930

Germany ■ 02103-29-12000

Hong Kong ■ 800 933 965

Ireland ■ 1800 555 049

Italy ■ 800-787980

Japan ■ 03-6890-7300

Korea (South) ■ 080-000-7145

Luxembourg ■ 8002 2076

Mexico ■ 01-800-7742-436

The Netherlands ■ 0800 0229592

Norway ■ 800-18859

Singapore ■ 1800-742-4368

Spain ■ 91-630-7050

Sweden ■ 020-790282

Switzerland ■ 055-254-22-11

UK ■ 01293-422-911

USA ■ 800-426-8157



Sample & Assay Technologies