RT² Profiler PCR Array (Rotor-Gene® Format) Mouse Inflammatory Cytokines & Receptors

Cat. no. 330231 PAMM-011ZR

For pathway expression analysis

Format	For use with the following real-time cyclers		
RT ² Profiler PCR Array,	Rotor-Gene Q, other Rotor-Gene cyclers		
Format R			

Description

The Mouse Inflammatory Cytokines & Receptors RT2 Profiler PCR Array profiles the expression of 84 key genes mediating the inflammatory response. Acute inflammation occurs in response to cell damage due to infection or injury. During this process, cellular and plasma derived factors encourage extravasation, the recruitment of circulating immune cells into the affected tissue. These immune cells in turn up-regulate the expression of inflammatory cytokines that recruit additional immune cells to mount an immune response to any invading organisms and to further promote and eventually resolve the inflammatory response. Chronic inflammation, or expression of these cytokines and receptors at low levels over long periods of time, promotes various pathological conditions including allergies and asthma, cardiovascular system (atherosclerosis), central nervous system disorders (Alzheimer's disease), fibrosis, and rheumatoid arthritis. This array contains genes involved in mediating immune cascade reactions during inflammation. The chemokines, cytokines, and interleukins involved in the inflammatory response are represented as well as their receptors. Profiling the expression of these inflammatory cytokine and receptor genes helps determine the state of and the mechanisms behind inflammation in your experimental model system. Using real-time PCR, you can easily and reliably analyze expression of a focused panel of genes related to inflammation with this array.

For further details, consult the RT² Profiler PCR Array Handbook.

Shipping and storage

RT² Profiler PCR Arrays in the Rotor-Gene format are shipped at ambient temperature, on dry ice, or blue ice packs depending on destination and accompanying products.

For long term storage, keep plates at -20°C.

Note: Ensure that you have the correct RT² Profiler PCR Array format for your real-time cycler (see table above).

Note: Open the package and store the products appropriately immediately on receipt.



Array layout

The 96 real-time assays in the Rotor-Gene format are located in wells 1–96 of the Rotor-Disc[™] (plate A1–A12=Rotor-Disc 1–12, plate B1–B12=Rotor-Disc 13–24, etc.). To maintain data analysis compatibility, wells 97–100 do not contain real-time assays but will contain master mix to account for weight balance.

Gene table: RT² Profiler PCR Array

Position	UniGene	GenBank	Symbol	Description	
A01	Mm.235137	NM_007926	Aimp1	Aminoacyl tRNA synthetase complex-interacting multifunctional protein 1	
A02	Mm.103205	NM 007553	Bmp2	Bone morphogenetic protein 2	
A03	Mm.1283	NM 011329	Ccl1	Chemokine (C-C motif) ligand 1	
A04	Mm.4686	NM 011330	Ccl11	Chemokine (C-C motif) ligand 11	
A05	Mm.867	NM 011331	Ccl12	Chemokine (C-C motif) ligand 12	
A06	Mm.41988	NM 011332	Ccl17	Chemokine (C-C motif) ligand 17	
A07	Mm.424740	NM 011888	Ccl19	Chemokine (C-C motif) ligand 19	
A08	Mm.290320	NM 011333	Ccl2	Chemokine (C-C motif) ligand 2	
A09	Mm.116739	NM 016960	Ccl20	Chemokine (C-C motif) ligand 20	
A10	Mm.12895	NM 009137	Ccl22	Chemokine (C-C motif) ligand 22	
A11	Mm.31505	NM 019577	Ccl24	Chemokine (C-C motif) ligand 24	
A12	Mm.1282	NM 011337	Ccl3	Chemokine (C-C motif) ligand 3	
B01	Mm.244263	NM 013652	Ccl4	Chemokine (C-C motif) ligand 4	
B02	Mm.284248	NM 013653	Ccl5	Chemokine (C-C motif) ligand 5	
B03	Mm.137	NM 009139	Ccl6	Chemokine (C-C motif) ligand 6	
B03	Mm.341574	NM 013654	Ccl7	Chemokine (C-C motif) ligand 7	
B05	Mm.42029	NM 021443	Ccl7	, , ,	
				Chemokine (C-C motif) ligand 8	
B06 B07	Mm.416125	NM_011338	Ccl9 Ccr1	Chemokine (C-C motif) ligand 9	
	Mm.274927	NM_009912		Chemokine (C-C motif) receptor 1	
B08	Mm.8021	NM_007721	Ccr10	Chemokine (C-C motif) receptor 10	
B09	Mm.6272	NM_009915	Ccr2	Chemokine (C-C motif) receptor 2	
B10	Mm.57050	NM_009914	Ccr3	Chemokine (C-C motif) receptor 3	
B11	Mm.1337	NM_009916	Ccr4	Chemokine (C-C motif) receptor 4	
B12	Mm.14302	NM_009917	Ccr5	Chemokine (C-C motif) receptor 5	
C01	Mm.8007	NM_009835	Ccr6	Chemokine (C-C motif) receptor 6	
C02	Mm.442098	NM_007720	Ccr8	Chemokine (C-C motif) receptor 8	
C03	Mm.4861	NM_011616	Cd40lg	CD40 ligand	
C04	Mm.795	NM_007778	Csf1	Colony stimulating factor 1 (macrophage)	
C05	Mm.4922	NM_009969	Csf2	Colony stimulating factor 2 (granulocyte-macrophage)	
C06	Mm.1238	NM_009971	Csf3	Colony stimulating factor 3 (granulocyte)	
C07	Mm.103711	NM_009142	Cx3cl1	Chemokine (C-X3-C motif) ligand 1	
C08	Mm.21013	NM_008176	Cxcl1	Chemokine (C-X-C motif) ligand 1	
C09	Mm.877	NM_021274	Cxcl10	Chemokine (C-X-C motif) ligand 10	
C10	Mm.131723	NM_019494	Cxcl11	Chemokine (C-X-C motif) ligand 11	
C11	Mm.303231	NM_021704	Cxcl12	Chemokine (C-X-C motif) ligand 12	
C12	Mm.10116	NM_018866	Cxcl13	Chemokine (C-X-C motif) ligand 13	
D01	Mm.64326	NM_011339	Cxcl15	Chemokine (C-X-C motif) ligand 15	
D02	Mm.4660	NM_009141	Cxcl5	Chemokine (C-X-C motif) ligand 5	
D03	Mm.766	NM_008599	Cxcl9	Chemokine (C-X-C motif) ligand 9	
D04	Mm.234466	NM_009909	Cxcr2	Chemokine (C-X-C motif) receptor 2	
D05	Mm.12876	NM_009910	Cxcr3	Chemokine (C-X-C motif) receptor 3	
D06	Mm.6246	NM_007551	Cxcr5	Chemokine (C-X-C motif) receptor 5	
D07	Mm.3355	NM_010177	Fasl	Fas ligand (TNF superfamily, member 6)	
D08	Mm.240327	NM_008337	Ifng	Interferon gamma	
D09	Mm.379327	NM_008348	II10ra	Interleukin 10 receptor, alpha	
D10	Mm.4154	NM_008349	II10rb	Interleukin 10 receptor, beta	
D11	Mm.35814	NM 008350	II11	Interleukin 11	
D12	Mm.1284	NM_008355	II13	Interleukin 13	
E01	Mm.4392	NM 008357	II15	Interleukin 15	
E02	Mm.10137	NM 010551	II16	Interleukin 16	
E03	Mm.5419	NM 010552	II17a	Interleukin 17A	
E04	Mm.59313	NM 019508	II17b	Interleukin 17B	
E05	Mm.222807	NM 145856	II176	Interleukin 17F	
E06	Mm.15534	NM 010554	II171	Interleukin 1 alpha	
E07	Mm.222830	NM 008361	II1b	Interleukin 1 beta	
LU/	141111.775090		IIID IIIrl		
E08	Mm.896	NM 008362		Interleukin 1 receptor, type I	

Position	UniGene	GenBank	Symbol	Description	
E10	Mm.157689	NM_021782	II21	Interleukin 21	
E11	Mm.222632	NM_145636	1127	Interleukin 27	
E12	Mm.35287	NM_008368	II2rb	Interleukin 2 receptor, beta chain	
F01	Mm.2923	NM_013563	Il2rg	Interleukin 2 receptor, gamma chain	
F02	Mm.983	NM_010556	II3	Interleukin 3	
F03	Mm.182359	NM_133775	1133	Interleukin 33	
F04	Mm.276360	NM_021283	II4	Interleukin 4	
F05	Mm.4461	NM_010558	II5	Interleukin 5	
F06	Mm.3448	NM_008370	Il5ra	Interleukin 5 receptor, alpha	
F07	Mm.2856	NM_010559	ll6ra	Interleukin 6 receptor, alpha	
F08	Mm.4364	NM_010560	llóst	Interleukin 6 signal transducer	
F09	Mm.3825	NM_008371	117	Interleukin 7	
F10	Mm.87787	NM_010735	Lta	Lymphotoxin A	
F11	Mm.1715	NM_008518	Ltb	Lymphotoxin B	
F12	Mm.2326	NM_010798	Mif	Macrophage migration inhibitory factor	
G01	Mm.202727	NM_021524	Nampt	Nicotinamide phosphoribosyltransferase	
G02	Mm.131422	NM_001013365	Osm	Oncostatin M	
G03	Mm.332490	NM_019932	Pf4	Platelet factor 4	
G04	Mm.288474	NM_009263	Spp1	Secreted phosphoprotein 1	
G05	Mm.1293	NM_013693	Tnf	Tumor necrosis factor	
G06	Mm.15383	NM_008764	Tnfrsf11b	Tumor necrosis factor receptor superfamily, member 11b (osteoprotegerin)	
G07	Mm.1062	NM_009425	Tnfsf10	Tumor necrosis factor (ligand) superfamily, member 10	
G08	Mm.249221	NM_011613	Tnfsf11	Tumor necrosis factor (ligand) superfamily, member 11	
G09	Mm.8983	NM_023517	Tnfsf13	Tumor necrosis factor (ligand) superfamily, member 13	
G10	Mm.28835	NM_033622	Tnfsf13b	Tumor necrosis factor (ligand) superfamily, member 13b	
G11	Mm.4994	NM_009452	Tnfsf4	Tumor necrosis factor (ligand) superfamily, member 4	
G12	Mm.282184	NM_009505	Vegfa	Vascular endothelial growth factor A	
H01	Mm.328431	NM_007393	Actb	Actin, beta	
H02	Mm.163	NM_009735	B2m	Beta-2 microglobulin	
H03	Mm.343110	NM_008084	Gapdh	Glyceraldehyde-3-phosphate dehydrogenase	
H04	Mm.3317	NM_010368	Gusb	Glucuronidase, beta	
H05	Mm.2180	NM_008302	Hsp90ab1	Heat shock protein 90 alpha (cytosolic), class B member 1	
H06	N/A	SA_00106	MGDC	Mouse Genomic DNA Contamination	
H07	N/A	SA_00104	RTC	Reverse Transcription Control	
H08	N/A	SA_00104	RTC	Reverse Transcription Control	
H09	N/A	SA_00104	RTC	Reverse Transcription Control	
H10	N/A	SA_00103	PPC	Positive PCR Control	
H11	N/A	SA_00103	PPC	Positive PCR Control	
H12	N/A	SA_00103	PPC	Positive PCR Control	

Related products

For optimal performance, RT² Profiler PCR Arrays should be used together with the RT² First Strand Kit for cDNA synthesis and RT² SYBR[®] Green qPCR Mastermixes for PCR.

Product	Contents	Cat. no.
RT ² First Strand Kit (12)	Enzymes and reagents for cDNA synthesis	330401
RT ² SYBR Green ROX [™] FAST Mastermix (2)*	For 2 x 96 assays in 96-well plates; suitable for use with the Rotor-Gene Q and other Rotor-Gene cyclers	330620

^{*} Larger kit sizes available; please inquire.

RT² Profiler PCR Array products are intended for molecular biology applications. These products are not intended for the diagnosis, prevention, or treatment of a disease.

For up-to-date licensing information and product-specific disclaimers, see the respective QIAGEN kit handbook or user manual. QIAGEN kit handbooks and user manuals are available at www.qiagen. com or can be requested from QIAGEN Technical Services or your local distributor.

Trademarks: QIAGEN®, Rotor-Gene®, Rotor-Disc™ (QIAGEN Group); ROX™ (Applera Corporation or its subsidiaries); SYBR® (Molecular Probes, Inc.).

1067688 03/2011 © 2011 QIAGEN, all rights reserved.

www.aiaaen.com Australia • 1-800-243-800 Austria • 0800/281010 Belgium • 0800-79612 Brazil • 0800-557779

Canada • 800-572-9613 China • 8621-3865-3865 Denmark ■ 80-885945 Finland • 0800-914416 France • 01-60-920-930 Germany • 02103-29-12000 Hong Kong • 800 933 965

Ireland • 1800 555 049 Italy • 800-787980 Japan ■ 03-6890-7300 Korea (South) • 080-000-7145 Luxembourg ■ 8002 2076 Mexico • 01-800-7742-436 The Netherlands • 0800 0229592 USA • 800-426-8157

Norway • 800-18859 Singapore ■ 1800-742-4368 Spain ■ 91-630-7050 Sweden • 020-790282 Switzerland ■ 055-254-22-11 UK • 01293-422-911

