RT² Profiler PCR Array (Rotor-Gene® Format) Mouse Signal Transduction PathwayFinder

Cat. no. 330231 PAMM-014ZR

For pathway expression analysis

| Format | For use with the following real-time cyclers |
|-------------------------------------|--|
| RT ² Profiler PCR Array, | Rotor-Gene Q, other Rotor-Gene cyclers |
| Format R | |

Description

The Mouse Signal Transduction PathwayFinder RT2 Profiler PCR Array profiles the expression of 84 key genes responsive to signal transduction pathway activation or inhibition. Cellular signaling forms a complex network of gene interactions involving multiple signal transduction pathways. Each pathway ultimately increases or decreases the expression of its target genes resulting in alteration of cellular processes. Changes in target gene expression suggest signaling pathway activation or inhibition. However, gene expression results in the same pathway vary widely, depending on model systems and experimental conditions. Therefore, multiple target genes from each pathway should be examined to ensure accurate signaling pathway identification in a variety of model systems. In addition, analyzing multiple pathways simultaneously tests for signaling pathway crosstalk. This array includes target genes for 10 commonly studied signal transduction pathways, including pathways important for developmental, immunological, metabolic, and stress-activated processes. Results obtained with this array can suggest pathways that are potentially activated or inhibited by an experimental stimulus for further follow-up studies. Using real-time PCR, research studies can easily and reliably analyze the expression of a focused panel of genes involved in cellular signaling with this array.

For further details, consult the RT² Profiler PCR Array Handbook.

Shipping and storage

RT² Profiler PCR Arrays in the Rotor-Gene format are shipped at ambient temperature, on dry ice, or blue ice packs depending on destination and accompanying products.

For long term storage, keep plates at -20°C.

Note: Ensure that you have the correct RT² Profiler PCR Array format for your real-time cycler (see table above).

Note: Open the package and store the products appropriately immediately on receipt.



Array layout

The 96 real-time assays in the Rotor-Gene format are located in wells 1–96 of the Rotor-Disc[™] (plate A1–A12=Rotor-Disc 1–12, plate B1–B12=Rotor-Disc 13–24, etc.). To maintain data analysis compatibility, wells 97–100 do not contain real-time assays but will contain master mix to account for weight balance.

Gene table: RT² Profiler PCR Array

| Position | UniGene | GenBank | Symbol | Description | |
|----------|-----------|--------------|---------|--|--|
| A01 | Mm.427252 | NM 001033606 | Acsl3 | Acyl-CoA synthetase long-chain family member 3 | |
| A02 | Mm.391337 | NM 019477 | Acsl4 | Acyl-CoA synthetase long-chain family member 4 | |
| A03 | Mm.292056 | NM 027976 | Acsl5 | Acyl-CoA synthetase long-chain family member 5 | |
| A04 | Mm.1408 | NM 009627 | Adm | Adrenomedullin | |
| A05 | Mm.250265 | NM 009709 | Arnt | Aryl hydrocarbon receptor nuclear translocator | |
| A06 | Mm.641 | NM_009716 | Atf4 | Activating transcription factor 4 | |
| A07 | Mm,71710 | NM 015732 | Axin2 | Axin2 | |
| A08 | Mm.19904 | NM 007527 | Bax | Bcl2-associated X protein | |
| A09 | Mm.7660 | NM 133234 | Bbc3 | BCL2 binding component 3 | |
| A10 | Mm.257460 | NM 009741 | Bcl2 | B-cell leukemia/lymphoma 2 | |
| A11 | Mm.425593 | NM 009742 | Bcl2a1a | B-cell leukemia/lymphoma 2 related protein A1a | |
| A12 | Mm.238213 | NM 009743 | Bcl2l1 | Bcl2-like 1 | |
| B01 | Mm.2026 | NM_007464 | Birc3 | Baculoviral IAP repeat-containing 3 | |
| B02 | Mm.103205 | NM 007553 | Bmp2 | Bone morphogenetic protein 2 | |
| B02 | Mm.6813 | NM_007554 | Bmp4 | | |
| B03 | Mm.392646 | _ | | Bone morphogenetic protein 4 | |
| | | NM_007570 | Btg2 | B-cell translocation gene 2, anti-proliferative | |
| B05 | Mm.283682 | NM_139305 | Car9 | Carbonic anhydrase 9 | |
| B06 | Mm.284248 | NM_013653 | Ccl5 | Chemokine (C-C motif) ligand 5 | |
| B07 | Mm.273049 | NM_007631 | Ccnd1 | Cyclin D1 | |
| B08 | Mm.333406 | NM_009829 | Ccnd2 | Cyclin D2 | |
| B09 | Mm.195663 | NM_007669 | Cdkn1a | Cyclin-dependent kinase inhibitor 1A (P21) | |
| B10 | Mm.2958 | NM_009875 | Cdkn1b | Cyclin-dependent kinase inhibitor 1B | |
| B11 | Mm.347407 | NM_007679 | Cebpd | CCAAT/enhancer binding protein (C/EBP), delta | |
| B12 | Mm.307620 | NM_009949 | Cpt2 | Carnitine palmitoyltransferase 2 | |
| C01 | Mm.795 | NM_007778 | Csf1 | Colony stimulating factor 1 (macrophage) | |
| C02 | Mm.240830 | NM_023118 | Dab2 | Disabled homolog 2 (Drosophila) | |
| C03 | Mm.8534 | NM_007912 | Egfr | Epidermal growth factor receptor | |
| C04 | Mm.182785 | NM_010128 | Emp1 | Epithelial membrane protein 1 | |
| C05 | Mm.349116 | NM_007942 | Еро | Erythropoietin | |
| C06 | Mm.22126 | NM_017399 | Fabp1 | Fatty acid binding protein 1, liver | |
| C07 | Mm.1626 | NM_007987 | Fas | Fas (TNF receptor superfamily member 6) | |
| C08 | Mm.1233 | NM_013517 | Fcer2a | Fc receptor, IgE, low affinity II, alpha polypeptide | |
| C09 | Mm.6215 | NM_010235 | Fosl1 | Fos-like antigen 1 | |
| C10 | Mm.1776 | NM_010239 | Fth1 | Ferritin heavy chain 1 | |
| C11 | Mm.72235 | NM_007836 | Gadd45a | Growth arrest and DNA-damage-inducible 45 alpha | |
| C12 | Mm.1360 | NM_008655 | Gadd45b | Growth arrest and DNA-damage-inducible 45 beta | |
| D01 | Mm.313866 | NM_008091 | Gata3 | GATA binding protein 3 | |
| D02 | Mm.485389 | NM 010295 | Gclc | Glutamate-cysteine ligase, catalytic subunit | |
| D03 | Mm.292676 | NM 008129 | Gclm | Glutamate-cysteine ligase, modifier subunit | |
| D04 | Mm.283573 | NM 010344 | Gsr | Glutathione reductase | |
| | | | | Homocysteine-inducible, endoplasmic reticulum stress-inducible, ubiquitin-like | |
| D05 | Mm.29151 | NM_022331 | Herpud1 | domain member 1 | |
| D06 | Mm.390859 | NM 008235 | Hes1 | Hairy and enhancer of split 1 (Drosophila) | |
| D07 | Mm.137268 | NM 010419 | Hes5 | Hairy and enhancer of split 5 (Drosophila) | |
| D08 | Mm.29581 | NM 010423 | Hey1 | Hairy/enhancer-of-split related with YRPW motif 1 | |
| D09 | Mm.103573 | NM 013904 | Hey2 | Hairy/enhancer-of-split related with YRPW motif 2 | |
| D10 | Mm.103615 | NM 013905 | Heyl | Hairy/enhancer-of-split related with YRPW motif-like | |
| D11 | Mm.276389 | NM 010442 | Hmox1 | Heme oxygenase (decycling) 1 | |
| D12 | Mm.435508 | NM 010493 | lcam1 | Intercellular adhesion molecule 1 | |
| E01 | Mm.444 | NM 010495 | ld1 | Inhibitor of DNA binding 1 | |
| E02 | Mm.240327 | NM_010493 | Ifng | · · | |
| E02 | | _ | Ifrd1 | Interferon gamma | |
| | Mm.168 | NM_013562 | | Interferon-related developmental regulator 1 | |
| E04 | Mm.105218 | NM_008390 | Irf1 | Interferon regulatory factor 1 | |
| E05 | Mm.22398 | NM_013822 | Jag1 | Jagged 1 | |
| E06 | Mm.29324 | NM_010699 | Ldha | Lactate dehydrogenase A | |
| E07 | Mm.12834 | NM_008494 | Lfng | LFNG O-fucosylpeptide 3-beta-N-acetylglucosaminyltransferase | |
| E08 | Mm.348025 | NM_029796 | Lrg1 | Leucine-rich alpha-2-glycoprotein 1 | |

| Position | UniGene | GenBank | Symbol | Description | |
|----------|-----------|-----------|----------|---|--|
| E09 | Mm.1639 | NM_008562 | Mcl1 | Myeloid cell leukemia sequence 1 | |
| E10 | Mm.4825 | NM_010810 | Mmp7 | Matrix metallopeptidase 7 | |
| E11 | Mm.2444 | NM_010849 | Мус | Myelocytomatosis oncogene | |
| E12 | Mm.290610 | NM_008714 | Notch1 | Notch gene homolog 1 (Drosophila) | |
| F01 | Mm.252 | NM_008706 | Nqo1 | NAD(P)H dehydrogenase, quinone 1 | |
| F02 | Mm.293626 | NM_138648 | Olr1 | Oxidized low density lipoprotein (lectin-like) receptor 1 | |
| F03 | Mm.7141 | NM_011045 | Pcna | Proliferating cell nuclear antigen | |
| F04 | Mm.328914 | NM_011145 | Ppard | Peroxisome proliferator activator receptor delta | |
| F05 | Mm.228798 | NM_008957 | Ptch1 | Patched homolog 1 | |
| F06 | Mm.273862 | NM_009029 | Rb1 | Retinoblastoma 1 | |
| F07 | Mm.250422 | NM_008871 | Serpine1 | Serine (or cysteine) peptidase inhibitor, clade E, member 1 | |
| F08 | Mm.330113 | NM_011989 | Slc27a4 | Solute carrier family 27 (fatty acid transporter), member 4 | |
| F09 | Mm.21002 | NM_011400 | Slc2a1 | Solute carrier family 2 (facilitated glucose transporter), member 1 | |
| F10 | Mm.3468 | NM_007707 | Socs3 | Suppressor of cytokine signaling 3 | |
| F11 | Mm.210815 | NM_009166 | Sorbs1 | Sorbin and SH3 domain containing 1 | |
| F12 | Mm.40828 | NM_011018 | Sqstm1 | Sequestosome 1 | |
| G01 | Mm.277406 | NM_009283 | Stat1 | Signal transducer and activator of transcription 1 | |
| G02 | Mm.1293 | NM_013693 | Tnf | Tumor necrosis factor | |
| G03 | Mm.1062 | NM_009425 | Tnfsf10 | Tumor necrosis factor (ligand) superfamily, member 10 | |
| G04 | Mm.260618 | NM_011660 | Txn1 | Thioredoxin 1 | |
| G05 | Mm.210155 | NM_015762 | Txnrd1 | Thioredoxin reductase 1 | |
| G06 | Mm.282184 | NM_009505 | Vegfa | Vascular endothelial growth factor A | |
| G07 | Mm.10222 | NM_018865 | Wisp1 | WNT1 inducible signaling pathway protein 1 | |
| G08 | Mm.1123 | NM_021279 | Wnt1 | Wingless-related MMTV integration site 1 | |
| G09 | Mm.10740 | NM_009520 | Wnt2b | Wingless related MMTV integration site 2b | |
| G10 | Mm.1367 | NM_009522 | Wnt3a | Wingless-related MMTV integration site 3A | |
| G11 | Mm.287544 | NM_009524 | Wnt5a | Wingless-related MMTV integration site 5A | |
| G12 | Mm.268282 | NM_009526 | Wnt6 | Wingless-related MMTV integration site 6 | |
| H01 | Mm.328431 | NM_007393 | Actb | Actin, beta | |
| H02 | Mm.163 | NM_009735 | B2m | Beta-2 microglobulin | |
| H03 | Mm.343110 | NM_008084 | Gapdh | Glyceraldehyde-3-phosphate dehydrogenase | |
| H04 | Mm.3317 | NM_010368 | Gusb | Glucuronidase, beta | |
| H05 | Mm.2180 | NM_008302 | Hsp90ab1 | Heat shock protein 90 alpha (cytosolic), class B member 1 | |
| H06 | N/A | SA_00106 | MGDC | Mouse Genomic DNA Contamination | |
| H07 | N/A | SA_00104 | RTC | Reverse Transcription Control | |
| H08 | N/A | SA_00104 | RTC | Reverse Transcription Control | |
| H09 | N/A | SA_00104 | RTC | Reverse Transcription Control | |
| H10 | N/A | SA_00103 | PPC | Positive PCR Control | |
| H11 | N/A | SA_00103 | PPC | Positive PCR Control | |
| H12 | N/A | SA_00103 | PPC | Positive PCR Control | |

Related products

For optimal performance, RT² Profiler PCR Arrays should be used together with the RT² First Strand Kit for cDNA synthesis and RT² SYBR[®] Green qPCR Mastermixes for PCR.

| Product | Contents | Cat. no. |
|--|--|----------|
| RT ² First Strand Kit (12) | Enzymes and reagents for cDNA synthesis | 330401 |
| RT ² SYBR Green ROX [™] FAST Mastermix (2)* | For 2 x 96 assays in 96-well plates; suitable for use with the Rotor-Gene Q and other Rotor-Gene cyclers | 330620 |

^{*} Larger kit sizes available; please inquire.

RT² Profiler PCR Array products are intended for molecular biology applications. These products are not intended for the diagnosis, prevention, or treatment of a disease.

For up-to-date licensing information and product-specific disclaimers, see the respective QIAGEN kit handbook or user manual. QIAGEN kit handbooks and user manuals are available at www.qiagen. com or can be requested from QIAGEN Technical Services or your local distributor.

Trademarks: QIAGEN®, Rotor-Gene®, Rotor-Disc™ (QIAGEN Group); ROX™ (Applera Corporation or its subsidiaries); SYBR® (Molecular Probes, Inc.).

1067688 03/2011 © 2011 QIAGEN, all rights reserved.

www.aiaaen.com Australia • 1-800-243-800 Austria • 0800/281010 Belgium • 0800-79612 Brazil • 0800-557779

Canada • 800-572-9613 China • 8621-3865-3865 Denmark ■ 80-885945 Finland • 0800-914416 France • 01-60-920-930 Germany • 02103-29-12000 Hong Kong • 800 933 965

Ireland • 1800 555 049 Italy • 800-787980 Japan ■ 03-6890-7300 Korea (South) • 080-000-7145 Luxembourg ■ 8002 2076 Mexico • 01-800-7742-436 The Netherlands • 0800 0229592 USA • 800-426-8157

Norway • 800-18859 Singapore ■ 1800-742-4368 Spain ■ 91-630-7050 Sweden • 020-790282 Switzerland ■ 055-254-22-11 UK • 01293-422-911

