

RT² Profiler PCR Array (96-Well Format and 384-Well [4 x 96] Format)

Mouse T Cell Anergy & Immune Tolerance

Cat. no. 330231 PAMM-074ZA

For pathway expression analysis

Format	For use with the following real-time cyclers
RT ² Profiler PCR Array, Format A	Applied Biosystems® models 5700, 7000, 7300, 7500, 7700, 7900HT, ViiA™ 7 (96-well block); Bio-Rad® models iCycler®, iQ™ 5, MyiQ™, MyiQ2; Bio-Rad/MJ Research Chromo4™; Eppendorf® Mastercycler® ep realplex models 2, 2s, 4, 4s; Stratagene® models Mx3005P®, Mx3000P®; Takara TP-800
RT ² Profiler PCR Array, Format C	Applied Biosystems models 7500 (Fast block), 7900HT (Fast block), StepOnePlus™, ViiA 7 (Fast block)
RT ² Profiler PCR Array, Format D	Bio-Rad CFX96™; Bio-Rad/MJ Research models DNA Engine Opticon®, DNA Engine Opticon 2; Stratagene Mx4000®
RT ² Profiler PCR Array, Format E	Applied Biosystems models 7900HT (384-well block), ViiA 7 (384-well block); Bio-Rad CFX384™
RT ² Profiler PCR Array, Format F	Roche® LightCycler® 480 (96-well block)
RT ² Profiler PCR Array, Format G	Roche LightCycler 480 (384-well block)
RT ² Profiler PCR Array, Format H	Fluidigm® BioMark™



Sample & Assay Technologies

Description

The Mouse T-cell Anergy and Immune Tolerance RT² Profiler PCR Array profiles the expression of 84 key genes involved in T-cell anergy and immune tolerance. T-cells' functional unresponsiveness, termed anergy, is a cell-intrinsic mechanism that contributes to peripheral self-tolerance. This array contains T-cell and B-cell regulators, cytokines and receptors, TNF superfamily members and their receptors, and transcriptional factors. Using real-time PCR, you can easily and reliably analyze expression of a focused panel of genes involved in T-cell anergy and immune tolerance with this array.

For further details, consult the *RT² Profiler PCR Array Handbook*.

Shipping and storage

RT² Profiler PCR Arrays in formats A, C, D, E, F, and G are shipped at ambient temperature, on dry ice, or blue ice packs depending on destination and accompanying products. RT² Profiler PCR Arrays in format H are shipped on dry ice or blue ice packs.

For long term storage, keep plates at –20°C.

Note: Ensure that you have the correct RT² Profiler PCR Array format for your real-time cycler (see table above).

Note: Open the package and store the products appropriately immediately on receipt.

Array layout (96-well)

For 384-well 4 x 96 PCR arrays, genes are present in a staggered format. Refer to the *RT² Profiler PCR Array Handbook* for layout.

	1	2	3	4	5	6	7	8	9	10	11	12
A	Btla	Cblb	Ccl3	Ccr4	Cd27	Cd28	Cd40	Cd40lg	Cd70	Cdk2	Cdk4	Cma1
B	Csf1	Csf2	Ctla4	Dgka	Dgkz	Egr2	Egr3	Eomes	Fas	Fasl	Fos	Foxp1
C	Foxp2	Foxp3	Gata3	Gzmb	Hdac9	Icam1	Icos	Ifng	Il10	Il10ra	Il13	Il15
D	Il17a	Il1a	Il2	Il2ra	Il2rb	Il31	Il4	Il5	Il6	Il7r	Ing4	Irf4
E	Itch	Itga1	Jak1	Jak3	Jun	Lat	Lep	Lgals3	Lta	Mef2a	Nfatc1	Nfatc2
F	Nfatc3	Nfkb1	Nhlh2	Notch1	Pdcd1	Pf1	Prkcc	Ptger2	Ptgs2	Rnf128	Sei1	Stat3
G	Stat6	Tbx21	Tgfb1	Tnfrsf10b	Tnfrsf14	Tnfrsf18	Tnfrsf4	Tnfrsf8	Tnfrsf9	Tnfrsf10	Tnfrsf14	Tnfrsf8
H	Actb	B2m	Gapdh	Gusb	Hsp90ab1	MGDC	RTC	RTC	RTC	PPC	PPC	PPC

Gene table: RT² Profiler PCR Array

Position	UniGene	GenBank	Symbol	Description
A01	Mm.38199	NM_177584	Btla	B and T lymphocyte associated
A02	Mm.328206	NM_001033238	Cblb	Casitas B-lineage lymphoma b
A03	Mm.1282	NM_011337	Ccl3	Chemokine (C-C motif) ligand 3
A04	Mm.1337	NM_009916	Ccr4	Chemokine (C-C motif) receptor 4
A05	Mm.367714	NM_001033126	Cd27	CD27 antigen
A06	Mm.255003	NM_007642	Cd28	CD28 antigen
A07	Mm.271833	NM_011611	Cd40	CD40 antigen
A08	Mm.4861	NM_011616	Cd40lg	CD40 ligand
A09	Mm.42228	NM_011617	Cd70	CD70 antigen
A10	Mm.111326	NM_016756	Cdk2	Cyclin-dependent kinase 2
A11	Mm.6839	NM_009870	Cdk4	Cyclin-dependent kinase 4
A12	Mm.1252	NM_010780	Cma1	Chymase 1, mast cell
B01	Mm.795	NM_007778	Csf1	Colony stimulating factor 1 (macrophage)
B02	Mm.4922	NM_009969	Csf2	Colony stimulating factor 2 (granulocyte-macrophage)
B03	Mm.390	NM_009843	Ctla4	Cytotoxic T-lymphocyte-associated protein 4
B04	Mm.291235	NM_016811	Dgka	Diacylglycerol kinase, alpha
B05	Mm.314923	NM_138306	Dgkz	Diacylglycerol kinase zeta
B06	Mm.290421	NM_010118	Egr2	Early growth response 2
B07	Mm.103737	NM_018781	Egr3	Early growth response 3
B08	Mm.200692	NM_010136	Eomes	Eomesodermin homolog (Xenopus laevis)
B09	Mm.1626	NM_007987	Fas	Fas (TNF receptor superfamily member 6)
B10	Mm.3355	NM_010177	Fasl	Fas ligand (TNF superfamily, member 6)
B11	Mm.246513	NM_010234	Fos	FBJ osteosarcoma oncogene
B12	Mm.234965	NM_053202	Foxp1	Forkhead box P1
C01	Mm.332919	NM_053242	Foxp2	Forkhead box P2
C02	Mm.182291	NM_054039	Foxp3	Forkhead box P3
C03	Mm.313866	NM_008091	Gata3	GATA binding protein 3
C04	Mm.14874	NM_013542	Gzmb	Granzyme B
C05	Mm.310551	NM_024124	Hdac9	Histone deacetylase 9
C06	Mm.435508	NM_010493	Icam1	Intercellular adhesion molecule 1
C07	Mm.42044	NM_017480	Icos	Inducible T-cell co-stimulator
C08	Mm.240327	NM_008337	Ifng	Interferon gamma
C09	Mm.874	NM_010548	Il10	Interleukin 10
C10	Mm.379327	NM_008348	Il10ra	Interleukin 10 receptor, alpha
C11	Mm.1284	NM_008355	Il13	Interleukin 13
C12	Mm.4392	NM_008357	Il15	Interleukin 15
D01	Mm.5419	NM_010552	Il17a	Interleukin 17A
D02	Mm.15534	NM_010554	Il1a	Interleukin 1 alpha
D03	Mm.14190	NM_008366	Il2	Interleukin 2
D04	Mm.915	NM_008367	Il2ra	Interleukin 2 receptor, alpha chain
D05	Mm.35287	NM_008368	Il2rb	Interleukin 2 receptor, beta chain
D06	Mm.75049	NM_029594	Il31	Interleukin 31
D07	Mm.276360	NM_021283	Il4	Interleukin 4
D08	Mm.4461	NM_010558	Il5	Interleukin 5
D09	Mm.1019	NM_031168	Il6	Interleukin 6

Position	UniGene	GenBank	Symbol	Description
D10	Mm.389	NM_008372	Il7r	Interleukin 7 receptor
D11	Mm.262547	NM_133345	Ing4	Inhibitor of growth family, member 4
D12	Mm.4677	NM_013674	Irf4	Interferon regulatory factor 4
E01	Mm.208286	NM_008395	Itch	Itchy, E3 ubiquitin protein ligase
E02	Mm.482186	NM_001033228	Itga1	Integrin alpha 1
E03	Mm.289657	NM_146145	Jak1	Janus kinase 1
E04	Mm.249645	NM_010589	Jak3	Janus kinase 3
E05	Mm.275071	NM_010591	Jun	Jun oncogene
E06	Mm.10280	NM_010689	Lat	Linker for activation of T cells
E07	Mm.277072	NM_008493	Lep	Leptin
E08	Mm.248615	NM_010705	Lgals3	Lectin, galactose binding, soluble 3
E09	Mm.87787	NM_010735	Lta	Lymphotoxin A
E10	Mm.132788	NM_001033713	Mef2a	Myocyte enhancer factor 2A
E11	Mm.329560	NM_016791	Nfatc1	Nuclear factor of activated T-cells, cytoplasmic, calcineurin-dependent 1
E12	Mm.116802	NM_010899	Nfatc2	Nuclear factor of activated T-cells, cytoplasmic, calcineurin-dependent 2
F01	Mm.383185	NM_010901	Nfatc3	Nuclear factor of activated T-cells, cytoplasmic, calcineurin-dependent 3
F02	Mm.256765	NM_008689	Nfkb1	Nuclear factor of kappa light polypeptide gene enhancer in B-cells 1, p105
F03	Mm.137286	NM_178777	Nhlh2	Nescent helix loop helix 2
F04	Mm.290610	NM_008714	Notch1	Notch gene homolog 1 (Drosophila)
F05	Mm.5024	NM_008798	Pdcd1	Programmed cell death 1
F06	Mm.240313	NM_011073	Prf1	Perforin 1 (pore forming protein)
F07	Mm.7980	NM_011102	Prkcc	Protein kinase C, gamma
F08	Mm.4630	NM_008964	Ptger2	Prostaglandin E receptor 2 (subtype EP2)
F09	Mm.292547	NM_011198	Ptgs2	Prostaglandin-endoperoxide synthase 2
F10	Mm.27764	NM_023270	Rnf128	Ring finger protein 128
F11	Mm.1461	NM_011346	Sell	Selectin, lymphocyte
F12	Mm.249934	NM_011486	Stat3	Signal transducer and activator of transcription 3
G01	Mm.121721	NM_009284	Stat6	Signal transducer and activator of transcription 6
G02	Mm.477879	NM_019507	Tbx21	T-box 21
G03	Mm.248380	NM_011577	Tgfb1	Transforming growth factor, beta 1
G04	Mm.193430	NM_020275	Tnfrsf10b	Tumor necrosis factor receptor superfamily, member 10b
G05	Mm.215147	NM_178931	Tnfrsf14	Tumor necrosis factor receptor superfamily, member 14 (herpesvirus entry mediator)
G06	Mm.482508	NM_009400	Tnfrsf18	Tumor necrosis factor receptor superfamily, member 18
G07	Mm.13885	NM_011659	Tnfrsf4	Tumor necrosis factor receptor superfamily, member 4
G08	Mm.12810	NM_009401	Tnfrsf8	Tumor necrosis factor receptor superfamily, member 8
G09	Mm.244187	NM_011612	Tnfrsf9	Tumor necrosis factor receptor superfamily, member 9
G10	Mm.1062	NM_009425	Tnfsf10	Tumor necrosis factor (ligand) superfamily, member 10
G11	Mm.483369	NM_019418	Tnfsf14	Tumor necrosis factor (ligand) superfamily, member 14
G12	Mm.4664	NM_009403	Tnfsf8	Tumor necrosis factor (ligand) superfamily, member 8
H01	Mm.328431	NM_007393	Actb	Actin, beta
H02	Mm.163	NM_009735	B2m	Beta-2 microglobulin
H03	Mm.343110	NM_008084	Gapdh	Glyceraldehyde-3-phosphate dehydrogenase
H04	Mm.3317	NM_010368	Gusb	Glucuronidase, beta
H05	Mm.2180	NM_008302	Hsp90ab1	Heat shock protein 90 alpha (cytosolic), class B member 1
H06	N/A	SA_00106	MGDC	Mouse Genomic DNA Contamination
H07	N/A	SA_00104	RTC	Reverse Transcription Control
H08	N/A	SA_00104	RTC	Reverse Transcription Control
H09	N/A	SA_00104	RTC	Reverse Transcription Control
H10	N/A	SA_00103	PPC	Positive PCR Control
H11	N/A	SA_00103	PPC	Positive PCR Control
H12	N/A	SA_00103	PPC	Positive PCR Control

Related products

For optimal performance, RT² Profiler PCR Arrays should be used together with the RT² First Strand Kit for cDNA synthesis and RT² SYBR[®] Green qPCR Mastermixes for PCR.

Product	Contents	Cat. no.
RT ² First Strand Kit (12)	Enzymes and reagents for cDNA synthesis	330401
RT ² SYBR Green qPCR Mastermix (2)*	For 2 x 96 assays in 96-well plates; suitable for use with real-time cyclers that do not require a reference dye, including: Bio-Rad models CFX96, CFX384, DNA Engine Opticon 2; Bio-Rad/MJ Research Chromo4; Roche LightCycler 480 (96-well and 384-well); all other cyclers	330500
RT ² SYBR Green ROX [™] qPCR Mastermix (2)*	For 2 x 96 assays in 96-well plates; suitable for use with the following real-time cyclers: Applied Biosystems models 5700, 7000, 7300, 7500 [Standard and FAST], 7700, 7900HT 96-well block [Standard and FAST] and 384-well block, StepOnePlus; Eppendorf Mastercycler ep realplex models 2, 2S, 4, 4S; Stratagene models Mx3000P, Mx3005P, Mx4000; Takara TP-800	330520
RT ² SYBR Green Fluor qPCR Mastermix (2)*	For 2 x 96 assays in 96-well plates; suitable for use with the following real-time cyclers: Bio-Rad models iCycler, iQ5, MyiQ, MyiQ2	330510

* Larger kit sizes available; please inquire.

RT² Profiler PCR Array products are intended for molecular biology applications. These products are not intended for the diagnosis, prevention, or treatment of a disease.

For up-to-date licensing information and product-specific disclaimers, see the respective QIAGEN kit handbook or user manual. QIAGEN kit handbooks and user manuals are available at www.qiagen.com or can be requested from QIAGEN Technical Services or your local distributor.

Trademarks: QIAGEN® (QIAGEN Group); Applied Biosystems®, ViiA™, StepOnePlus™, ROX™ (Applied Biosystems Corporation or its subsidiaries); Bio-Rad®, iCycler®, iQ™, MyiQ™, Chromo4™, CFX96™, DNA Engine Opticon®, CFX384™ (Bio-Rad Laboratories, Inc.); Stratagene®, Mx3005P®, Mx3000P®, Mx4000® (Stratagene); Eppendorf®, Mastercycler® (Eppendorf AG); Roche®, LightCycler® (Roche Group); Fluidigm® BioMark™ (Fluidigm Corporation); SYBR® (Molecular Probes, Inc.).

1066029 03/2011 © 2011 QIAGEN, all rights reserved.

www.qiagen.com

Canada ■ 800-572-9613

China ■ 8621-3865-3865

Denmark ■ 80-885945

Finland ■ 0800-914416

France ■ 01-60-920-930

Germany ■ 02103-29-12000

Hong Kong ■ 800 933 965

Ireland ■ 1800 555 049

Italy ■ 800-787980

Japan ■ 03-6890-7300

Korea (South) ■ 080-000-7145

Luxembourg ■ 8002 2076

Mexico ■ 01-800-7742-436

The Netherlands ■ 0800 0229592

Norway ■ 800-18859

Singapore ■ 1800-742-4368

Spain ■ 91-630-7050

Sweden ■ 020-790282

Switzerland ■ 055-254-22-11

UK ■ 01293-422-911

USA ■ 800-426-8157

Australia ■ 1-800-243-800

Austria ■ 0800/281010

Belgium ■ 0800-79612

Brazil ■ 0800-557779



Sample & Assay Technologies