

# RT<sup>2</sup> Profiler PCR Array (96-Well Format and 384-Well [4 x 96] Format)

## Mouse Female Infertility

Cat. no. 330231 PAMM-164ZA

For pathway expression analysis

Format	For use with the following real-time cyclers
RT <sup>2</sup> Profiler PCR Array, Format A	Applied Biosystems® models 5700, 7000, 7300, 7500, 7700, 7900HT, ViiA™ 7 (96-well block); Bio-Rad® models iCycler®, iQ™ 5, MyiQ™, MyiQ2; Bio-Rad/MJ Research Chromo4™; Eppendorf® Mastercycler® ep realplex models 2, 2s, 4, 4s; Stratagene® models Mx3005P®, Mx3000P®; Takara TP-800
RT <sup>2</sup> Profiler PCR Array, Format C	Applied Biosystems models 7500 (Fast block), 7900HT (Fast block), StepOnePlus™, ViiA 7 (Fast block)
RT <sup>2</sup> Profiler PCR Array, Format D	Bio-Rad CFX96™; Bio-Rad/MJ Research models DNA Engine Opticon®, DNA Engine Opticon 2; Stratagene Mx4000®
RT <sup>2</sup> Profiler PCR Array, Format E	Applied Biosystems models 7900HT (384-well block), ViiA 7 (384-well block); Bio-Rad CFX384™
RT <sup>2</sup> Profiler PCR Array, Format F	Roche® LightCycler® 480 (96-well block)
RT <sup>2</sup> Profiler PCR Array, Format G	Roche LightCycler 480 (384-well block)
RT <sup>2</sup> Profiler PCR Array, Format H	Fluidigm® BioMark™



Sample & Assay Technologies

## Description

The Mouse Female Infertility RT<sup>2</sup> Profiler PCR Array profiles the expression of 84 key genes involved in endometrial changes in early pregnancy. During each human menstrual cycle, the uterine endometrial layer prepares for a pregnancy while one ovary releases a mature oocyte. If the oocyte is fertilized, it will begin to divide as it travels through the fallopian tube and into the uterus. The endometrium must undergo significant gene expression changes, required for successful embryonic implantation. These changes involve signaling pathways such as interleukin-1, WNT, and prostaglandins, as well as biological processes such as apoptosis, cell cycle regulation, and leukocyte migration into the receptive endometrium. The endometrium is only receptive to the embryo for 3–5 days per menstrual cycle, called the window of implantation. A large number of couples suffer from infertility, although there are now many assisted reproductive technologies that can achieve a viable pregnancy. While these technologies, such as in vitro fertilization (IVF), can identify the most promising embryos, more than half of IVF embryos fail to implant. A shorter or nonexistent window of implantation may explain this phenomenon for some infertile women. Many microarray studies have been published identifying endometrial gene expression changes during the window of implantation, as well as a comparison of fertile and infertile women. Additional studies are still required to identify the underlying molecular mechanisms of this complex process, potentially determined by monitoring expression of the key genes included on this array. A set of controls present on each array enables data analysis using the  $\Delta\Delta\text{CT}$  method of relative quantification, assessment of reverse transcription performance, genomic DNA contamination, and PCR performance. Using real-time PCR, research studies can easily and reliably analyze the expression of a focused panel of genes involved in embryonic implantation and female infertility with this array.

For further details, consult the *RT<sup>2</sup> Profiler PCR Array Handbook*.

## Shipping and storage

RT<sup>2</sup> Profiler PCR Arrays in formats A, C, D, E, F, and G are shipped at ambient temperature, on dry ice, or blue ice packs depending on destination and accompanying products. RT<sup>2</sup> Profiler PCR Arrays in format H are shipped on dry ice or blue ice packs.

For long term storage, keep plates at  $-20^{\circ}\text{C}$ .

**Note:** Ensure that you have the correct RT<sup>2</sup> Profiler PCR Array format for your real-time cycler (see table above).

**Note:** Open the package and store the products appropriately immediately on receipt.

---

## Array layout (96-well)

For 384-well 4 x 96 PCR arrays, genes are present in a staggered format. Refer to the *RT<sup>2</sup> Profiler PCR Array Handbook* for layout.

	1	2	3	4	5	6	7	8	9	10	11	12
<b>A</b>	Akt1	Anxa2	Apod	Ar	Areg	Bax	Bcl2	C2	C3	Calca	Casp3	Ccl5
<b>B</b>	Ccnb1	Cd55	Cdh1	Cfd	Cldn4	Comp	Crabp2	Csf1	Ctnnb1	Cxcl12	Dkk1	Egf
<b>C</b>	Egfr	Esr1	Esr2	F3	Fbn1	Fn1	Gadd45a	Gast	Gdf15	Gpx3	Hbegf	Hoxa10
<b>D</b>	Hoxa11	Icam1	Igf1	Igfbp1	Il11	Il15	Il1a	Il1b	Il1r1	Il6	Ilgf4	Ilgf5
<b>E</b>	Ilgb3	Kdr	Lama1	Lamc2	Lep	Lif	Lifr	Maoa	Mid1	Mki67	Mmp2	Mmp7
<b>F</b>	Mmp9	Msx1	Muc1	Olfm1	Pcna	Pgf	Pgr	Pri	Ptgs1	Ptgs2	Sell	Sfrp4
<b>G</b>	Sod1	Spp1	Stat3	Stmn1	Tgfb1	Timp1	Tnf	Tnfrsf10b	Trp53	Vcam1	Vegfa	Wnt2
<b>H</b>	Actb	B2m	Gapdh	Gusb	Hsp90ab1	MGDC	RTC	RTC	RTC	PPC	PPC	PPC

## Gene table: RT<sup>2</sup> Profiler PCR Array

Position	UniGene	GenBank	Symbol	Description
A01	Mm.6645	NM_009652	Akt1	Thymoma viral proto-oncogene 1
A02	Mm.238343	NM_007585	Anxa2	Annexin A2
A03	Mm.2082	NM_007470	Apod	Apolipoprotein D
A04	Mm.39005	NM_013476	Ar	Androgen receptor
A05	Mm.8039	NM_009704	Areg	Amphiregulin
A06	Mm.19904	NM_007527	Bax	Bcl2-associated X protein
A07	Mm.257460	NM_009741	Bcl2	B-cell leukemia/lymphoma 2
A08	Mm.283217	NM_013484	C2	Complement component 2 (within H-2S)
A09	Mm.19131	NM_009778	C3	Complement component 3
A10	Mm.4361	NM_007587	Calca	Calcitonin/calcitonin-related polypeptide, alpha
A11	Mm.34405	NM_009810	Casp3	Caspase 3
A12	Mm.284248	NM_013653	Ccl5	Chemokine (C-C motif) ligand 5
B01	Mm.260114	NM_172301	Ccnb1	Cyclin B1
B02	Mm.101591	NM_010016	Cd55	CD55 antigen
B03	Mm.35605	NM_009864	Cdh1	Cadherin 1
B04	Mm.4407	NM_013459	Cfd	Complement factor D (adipsin)
B05	Mm.7339	NM_009903	Cldn4	Claudin 4
B06	Mm.45071	NM_016685	Comp	Cartilage oligomeric matrix protein
B07	Mm.4757	NM_007759	Crabp2	Cellular retinoic acid binding protein II
B08	Mm.795	NM_007778	Csf1	Colony stimulating factor 1 (macrophage)
B09	Mm.291928	NM_007614	Ctnnb1	Catenin (cadherin associated protein), beta 1
B10	Mm.303231	NM_021704	Cxcl12	Chemokine (C-X-C motif) ligand 12
B11	Mm.214717	NM_010051	Dkk1	Dickkopf homolog 1 (Xenopus laevis)
B12	Mm.252481	NM_010113	Egf	Epidermal growth factor
C01	Mm.8534	NM_007912	Egfr	Epidermal growth factor receptor
C02	Mm.9213	NM_007956	Esr1	Estrogen receptor 1 (alpha)
C03	Mm.2561	NM_010157	Esr2	Estrogen receptor 2 (beta)
C04	Mm.273188	NM_010171	F3	Coagulation factor III
C05	Mm.271644	NM_007993	Fbn1	Fibrillin 1
C06	Mm.193099	NM_010233	Fn1	Fibronectin 1
C07	Mm.72235	NM_007836	Gadd45a	Growth arrest and DNA-damage-inducible 45 alpha
C08	Mm.4767	NM_010257	Gast	Gastrin
C09	Mm.31325	NM_011819	Gdf15	Growth differentiation factor 15
C10	Mm.200916	NM_008161	Gpx3	Glutathione peroxidase 3
C11	Mm.289681	NM_010415	Hbegf	Heparin-binding EGF-like growth factor
C12	Mm.5	NM_008263	Hoxa10	Homeobox A10
D01	Mm.26954	NM_010450	Hoxa11	Homeobox A11
D02	Mm.435508	NM_010493	Icam1	Intercellular adhesion molecule 1
D03	Mm.268521	NM_010512	Igf1	Insulin-like growth factor 1
D04	Mm.21300	NM_008341	Igfbp1	Insulin-like growth factor binding protein 1
D05	Mm.35814	NM_008350	Il11	Interleukin 11
D06	Mm.4392	NM_008357	Il15	Interleukin 15
D07	Mm.15534	NM_010554	Il1a	Interleukin 1 alpha
D08	Mm.222830	NM_008361	Il1b	Interleukin 1 beta
D09	Mm.896	NM_008362	Il1r1	Interleukin 1 receptor, type I

Position	UniGene	GenBank	Symbol	Description
D10	Mm.1019	NM_031168	Il6	Interleukin 6
D11	Mm.31903	NM_010576	Itga4	Integrin alpha 4
D12	Mm.227	NM_008402	Itgav	Integrin alpha V
E01	Mm.87150	NM_016780	Itgb3	Integrin beta 3
E02	Mm.285	NM_010612	Kdr	Kinase insert domain protein receptor
E03	Mm.303386	NM_008480	Lama1	Laminin, alpha 1
E04	Mm.329272	NM_008485	Lamc2	Laminin, gamma 2
E05	Mm.277072	NM_008493	Lep	Leptin
E06	Mm.4964	NM_008501	Lif	Leukemia inhibitory factor
E07	Mm.149720	NM_013584	Lifr	Leukemia inhibitory factor receptor
E08	Mm.21108	NM_173740	Maoa	Monoamine oxidase A
E09	Mm.34441	NM_010797	Mid1	Midline 1
E10	Mm.4078	NM_001081117	Mki67	Antigen identified by monoclonal antibody Ki 67
E11	Mm.29564	NM_008610	Mmp2	Matrix metalloproteinase 2
E12	Mm.4825	NM_010810	Mmp7	Matrix metalloproteinase 7
F01	Mm.4406	NM_013599	Mmp9	Matrix metalloproteinase 9
F02	Mm.256509	NM_010835	Msx1	Homeobox, msh-like 1
F03	Mm.16193	NM_013605	Muc1	Mucin 1, transmembrane
F04	Mm.43278	NM_019498	Olfm1	Olfactomedin 1
F05	Mm.7141	NM_011045	Pcna	Proliferating cell nuclear antigen
F06	Mm.4809	NM_008827	Pgf	Placental growth factor
F07	Mm.12798	NM_008829	Pgr	Progesterone receptor
F08	Mm.1270	NM_011164	Prl	Prolactin
F09	Mm.275434	NM_008969	Ptgs1	Prostaglandin-endoperoxide synthase 1
F10	Mm.292547	NM_011198	Ptgs2	Prostaglandin-endoperoxide synthase 2
F11	Mm.1461	NM_011346	Sell	Selectin, lymphocyte
F12	Mm.42095	NM_016687	Sfrp4	Secreted frizzled-related protein 4
G01	Mm.276325	NM_011434	Sod1	Superoxide dismutase 1, soluble
G02	Mm.288474	NM_009263	Spp1	Secreted phosphoprotein 1
G03	Mm.249934	NM_011486	Stat3	Signal transducer and activator of transcription 3
G04	Mm.378957	NM_019641	Stmn1	Stathmin 1
G05	Mm.248380	NM_011577	Tgfb1	Transforming growth factor, beta 1
G06	Mm.8245	NM_011593	Timp1	Tissue inhibitor of metalloproteinase 1
G07	Mm.1293	NM_013693	Tnf	Tumor necrosis factor
G08	Mm.193430	NM_020275	Tnfrsf10b	Tumor necrosis factor receptor superfamily, member 10b
G09	Mm.222	NM_011640	Trp53	Transformation related protein 53
G10	Mm.76649	NM_011693	Vcam1	Vascular cell adhesion molecule 1
G11	Mm.282184	NM_009505	Vegfa	Vascular endothelial growth factor A
G12	Mm.33653	NM_023653	Wnt2	Wingless-related MMTV integration site 2
H01	Mm.328431	NM_007393	Actb	Actin, beta
H02	Mm.163	NM_009735	B2m	Beta-2 microglobulin
H03	Mm.343110	NM_008084	Gapdh	Glyceraldehyde-3-phosphate dehydrogenase
H04	Mm.3317	NM_010368	Gusb	Glucuronidase, beta
H05	Mm.2180	NM_008302	Hsp90ab1	Heat shock protein 90 alpha (cytosolic), class B member 1
H06	N/A	SA_00106	MGDC	Mouse Genomic DNA Contamination
H07	N/A	SA_00104	RTC	Reverse Transcription Control
H08	N/A	SA_00104	RTC	Reverse Transcription Control
H09	N/A	SA_00104	RTC	Reverse Transcription Control
H10	N/A	SA_00103	PPC	Positive PCR Control
H11	N/A	SA_00103	PPC	Positive PCR Control
H12	N/A	SA_00103	PPC	Positive PCR Control

## Related products

For optimal performance, RT<sup>2</sup> Profiler PCR Arrays should be used together with the RT<sup>2</sup> First Strand Kit for cDNA synthesis and RT<sup>2</sup> SYBR<sup>®</sup> Green qPCR Mastermixes for PCR.

Product	Contents	Cat. no.
RT <sup>2</sup> First Strand Kit (12)	Enzymes and reagents for cDNA synthesis	330401
RT <sup>2</sup> SYBR Green qPCR Mastermix (2)*	For 2 x 96 assays in 96-well plates; suitable for use with real-time cyclers that do not require a reference dye, including: Bio-Rad models CFX96, CFX384, DNA Engine Opticon 2; Bio-Rad/MJ Research Chromo4; Roche LightCycler 480 (96-well and 384-well); all other cyclers	330500
RT <sup>2</sup> SYBR Green ROX™ qPCR Mastermix (2)*	For 2 x 96 assays in 96-well plates; suitable for use with the following real-time cyclers: Applied Biosystems models 5700, 7000, 7300, 7500 [Standard and FAST], 7700, 7900HT 96-well block [Standard and FAST] and 384-well block, StepOnePlus; Eppendorf Mastercycler ep realplex models 2, 2S, 4, 4S; Stratagene models Mx3000P, Mx3005P, Mx4000; Takara TP-800	330520
RT <sup>2</sup> SYBR Green Fluor qPCR Mastermix (2)*	For 2 x 96 assays in 96-well plates; suitable for use with the following real-time cyclers: Bio-Rad models iCycler, iQ5, MyiQ, MyiQ2	330510

\* Larger kit sizes available; please inquire.

RT<sup>2</sup> Profiler PCR Array products are intended for molecular biology applications. These products are not intended for the diagnosis, prevention, or treatment of a disease.

For up-to-date licensing information and product-specific disclaimers, see the respective QIAGEN kit handbook or user manual. QIAGEN kit handbooks and user manuals are available at [www.qiagen.com](http://www.qiagen.com) or can be requested from QIAGEN Technical Services or your local distributor.

Trademarks: QIAGEN® (QIAGEN Group); Applied Biosystems®, ViiA™, StepOnePlus™, ROX™ (Applied Biosystems Corporation or its subsidiaries); Bio-Rad®, iCycler®, iQ™, MyiQ™, Chromo4™, CFX96™, DNA Engine Opticon®, CFX384™ (Bio-Rad Laboratories, Inc.); Stratagene®, Mx3005P®, Mx3000P®, Mx4000® (Stratagene); Eppendorf®, Mastercycler® (Eppendorf AG); Roche®, LightCycler® (Roche Group); Fluidigm® BioMark™ (Fluidigm Corporation); SYBR® (Molecular Probes, Inc.).

1066029 03/2011 © 2011 QIAGEN, all rights reserved.

[www.qiagen.com](http://www.qiagen.com)

Canada ■ 800-572-9613

Ireland ■ 1800 555 049

Norway ■ 800-18859

China ■ 8621-3865-3865

Italy ■ 800-787980

Singapore ■ 1800-742-4368

Denmark ■ 80-885945

Japan ■ 03-6890-7300

Spain ■ 91-630-7050

Australia ■ 1-800-243-800

Finland ■ 0800-914416

Korea (South) ■ 080-000-7145

Sweden ■ 020-790282

Austria ■ 0800/281010

France ■ 01-60-920-930

Luxembourg ■ 8002 2076

Switzerland ■ 055-254-22-11

Belgium ■ 0800-79612

Germany ■ 02103-29-12000

Mexico ■ 01-800-7742-436

UK ■ 01293-422-911

Brazil ■ 0800-557779

Hong Kong ■ 800 933 965

The Netherlands ■ 0800 0229592

USA ■ 800-426-8157



Sample & Assay Technologies