RT² Profiler PCR Array (96-Well Format and 384-Well [4 x 96] Format)

Rhesus Macaque Apoptosis

Cat. no. 330231 PAQQ-012ZA

For pathway expression analysis

Format	For use with the following real-time cyclers
RT ² Profiler PCR Array, Format A	Applied Biosystems® models 5700, 7000, 7300, 7500, 7700, 7900HT, ViiA™ 7 (96-well block); Bio-Rad® models iCycler®, iQ™5, MyiQ™, MyiQ2; Bio-Rad/MJ Research Chromo4™; Eppendorf® Mastercycler® ep realplex models 2, 2s, 4, 4s; Stratagene® models Mx3005P®, Mx3000P®; Takara TP-800
RT ² Profiler PCR Array, Format C	Applied Biosystems models 7500 (Fast block), 7900HT (Fast block), StepOnePlus™, ViiA 7 (Fast block)
RT ² Profiler PCR Array, Format D	Bio-Rad CFX96™; Bio-Rad/MJ Research models DNA Engine Opticon®, DNA Engine Opticon 2; Stratagene Mx4000®
RT ² Profiler PCR Array, Format E	Applied Biosystems models 7900HT (384-well block), ViiA 7 (384-well block); Bio-Rad CFX384™
RT ² Profiler PCR Array, Format F	Roche® LightCycler® 480 (96-well block)
RT ² Profiler PCR Array, Format G	Roche LightCycler 480 (384-well block)
RT ² Profiler PCR Array, Format H	Fluidigm [®] BioMark™



Description

The Rhesus Macaque Apoptosis RT2 Profiler PCR Array profiles the expression of 84 key genes involved in programmed cell death. Apoptosis plays a critical role in normal biological processes requiring cell removal, including differentiation, development, and homeostasis. Stress responses (such as heat shock, ischemia, unfolded proteins, and viral infection) cause badly damaged cells to undergo apoptosis. In cell culture, growth factor withdrawal and many known experimental compounds have a similar effect. An acquired defect in apoptosis activation often leads to uncontrolled cell growth, oncogenesis, and cancer. Ligand-bound tumor necrosis factor (TNF) receptors initiate apoptosis by recruiting FADD and other death domain-containing adaptor proteins that then recruit and activate caspases. Environmental stresses trigger BCL2 protein oligomerization and insertion into the mitochondrial membrane, releasing APAF1 and other CARD family members that also oligomerize to recruit and activate caspases. Caspases promote a proteolysis cascade that degrades cellular protein targets, while the IAP protein family directly inhibits caspases. This array includes TNF ligands and their receptors, members of the Bcl-2, caspase, IAP, TRAF, CARD, death domain, death-effector domain, and CIDE families, as well as genes involved in the p53 and DNA damage pathways. Monitoring the expression of these genes helps determine the mechanisms behind programmed cell death in model systems and the propensity of a cell type to undergo apoptosis normally. Using real-time PCR, research studies can easily and reliably analyze expression of a focused panel of genes related to apoptosis with this array.

For further details, consult the RT² Profiler PCR Array Handbook.

Shipping and storage

RT² Profiler PCR Arrays in formats A, C, D, E, F, and G are shipped at ambient temperature, on dry ice, or blue ice packs depending on destination and accompanying products. RT² Profiler PCR Arrays in format H are shipped on dry ice or blue ice packs.

For long term storage, keep plates at -20°C.

Note: Ensure that you have the correct RT² Profiler PCR Array format for your real-time cycler (see table above).

Note: Open the package and store the products appropriately immediately on receipt.

Array layout (96-well)

For 384-well 4 x 96 PCR arrays, genes are present in a staggered format. Refer to the RT^2 Profiler PCR Array Handbook for layout.

	1	2	3	4	5	6	7	8	9	10	11	12
А	ABL1	AKT1	APAF1	BAD	BAG1	BAG3	BAG4	BAK1	BAX	BCL2A1	BCL2L1	BCL2L10
В	BCL2L11	BDNF	BFAR	BIRC2	BIRC3	BIRC5	BIRC6	BNIP1	BNIP2	BNIP3L	CARD8	CASP1
С	CASP10	CASP14	CASP2	CASP3	CASP4	CASP5	CASP7	CASP8	CASP9	CD27	CD40	CD40LG
D	CD70	CDKN1A	CFLAR	CIDEA	CRADD	DAPK1	DFFA	FAS	FASLG	GADD45A	IGF1R	IL10
E	IL2	LOC1004261	LOC693554	LOC695164	LOC698160	LOC698788	LOC707407	LOC708606	LOC714869	LOC716650	LTA	LTBR
F	MAPK1	MCL1	MYC	NAIP	NOD1	NOL3	PRKCA	PYCARD	RIPK2	TNF	TNFRSF10A	TNFRSF11B
G	TNFRSF1A	TNFRSF21	TNFRSF25	TNFRSF9	TNFSF10	TNFSF8	TP53	TP53BP2	TRAF3	TRAF4	TRAF5	XIAP
н	ACTB	B2M	GAPDH	LOC709186	RPL13A	QGDC	RTC	RTC	RTC	PPC	PPC	PPC

Gene table: RT² Profiler PCR Array

Position	UniGene	GenBank	Symbol	Description
A01	N/A	XM_001118598	ABL1	C-abl oncogene 1, non-receptor tyrosine kinase
A02	Mmu.1599	XM_001085265	AKT1	V-akt murine thymoma viral oncogene homolog 1
A03	Mmu.15956	XM_001086840	APAF1	Apoptotic peptidase activating factor 1
A04	Mmu.18791	XM_001115090	BAD	BCL2-associated agonist of cell death
A05	Mmu.2086	XM_001090384	BAG1	BCL2-associated athanogene
A06	N/A	XM_001104160	BAG3	BCL2-associated athanogene 3
A07	Mmu.14159	XM_001092290	BAG4	BCL2-associated athanogene 4
A08	Mmu.11738	XM_001090570	BAK1	BCL2-antagonist/killer 1
A09	N/A	XM_001112353	BAX	BCL2-associated X protein
A10	Mmu.16575	XM_001109281	BCL2A1	BCL2-related protein A1
A11	Mmu.16103	XM 001110285	BCL2L1	BCL2-like 1
A12	N/A	XM 001085850	BCL2L10	BCL2-like 10 (apoptosis facilitator)
B01	Mmu.14594	XM 001086237	BCL2L11	BCL2-like 11 (apoptosis facilitator)
B02	Mmu.3380	XM 001089568	BDNF	Brain-derived neurotrophic factor
B03	Mmu.12798	XM 001107948	BFAR	Bifunctional apoptosis regulator
B04	Mmu.1906	XM 001097089	BIRC2	Baculoviral IAP repeat containing 2
B05	Mmu.2837	XM 001096429	BIRC3	Baculoviral IAP repeat containing 3
B06	N/A	XM 001108164	BIRC5	Baculoviral IAP repeat containing 5
B07	Mmu.9645	XM 002799203	BIRC6	Baculoviral IAP repeat-containing 6
B08	Mmu.16357	XM 001096579	BNIP1	BCL2/adenovirus E1B 19kDa interacting protein 1
B09	Mmu.16319	XM 001100604	BNIP2	BCL2/adenovirus E1B 19kDa interacting protein 2
B10	Mmu.4295	NM 001037284	BNIP3L	BCL2/adenovirus E1B 19kDa interacting protein 3-like
B11	N/A	XM 001111494	CARD8	Caspase recruitment domain family, member 8
B12	Mmu.11508	XM 001094943	CASP1	Caspase 1, apoptosis-related cysteine peptidase (interleukin 1, beta, convertase)
C01	Mmu.15917	XM 001097804	CASP10	Caspase 10, apoptosis-related cysteine peptidase
C02	N/A	XM 001111598	CASP14	Caspase 14, apoptosis-related cysteine peptidase
C03	N/A	XM 001087704	CASP2	Caspase-2-like
C04	Mmu,1947	XM 001083160	CASP3	Caspase 3, apoptosis-related cysteine peptidase
C05	Mmu.12625	XM 001100287	CASP4	Caspase 4, apoptosis-related cysteine peptidase
C06	N/A	XM 001100375	CASP5	Caspase 5, apoptosis-related cysteine peptidase
C07	Mmu.11428	XM 001090833	CASP7	Caspase 7, apoptosis-related cysteine peptidase
C08	N/A	XM 001091080	CASP8	Caspase 8, apoptosis-related cysteine peptidase
C09	N/A	XM 001082859	CASP9	Caspase 9, apoptosis-related cysteine peptidase
C10	Mmu.1817	XM 001104337	CD27	CD27 antigen-like
C11	N/A	XM 001104333	CD40	CD40 molecule, TNF receptor superfamily member 5
C12	Mmu.3581	NM 001032839	CD40LG	CD40 ligand
D01	N/A	XM 001088935	CD70	CD70 antigen-like
D02	Mmu.4393	NM 001194722	CDKN1A	Cyclin-dependent kinase inhibitor 1A (p21, Cip1)
D03	Mmu.3649	XM 001096860	CFLAR	CASP8 and FADD-like apoptosis regulator
D03	N/A	XM 001094509	CIDEA	Cell death-inducing DFFA-like effector a
D05	Mmu.11725	XM_001074307 XM_001105475	CRADD	CASP2 and RIPK1 domain containing adaptor with death domain
D05	Mmu.149	XM_001103473	DAPK1	Death-associated protein kinase 1
D07	N/A	XM_001084783	DFFA	DNA fragmentation factor, 45kDa, alpha polypeptide
D07	IN/A	741 UUTTUZ 193	אווע	Distribution racion, 45kDa, alpha polypepilde

Position	UniGene	GenBank	Symbol	Description
D08	Mmu.649	NM_001032933	FAS	Fas (TNF receptor superfamily, member 6)
D09	Mmu.3579	NM 001032838	FASLG	Fas ligand (TNF superfamily, member 6)
D10	Mmu.10930	XM_001095413	GADD45A	Growth arrest and DNA-damage-inducible, alpha
D11	N/A	XM_001100407	IGF1R	Insulin-like growth factor 1 receptor
D12	Mmu.3374	NM 001044727	IL10	Interleukin 10
E01	Mmu.3415	NM 001047130	IL2	Interleukin 2
E02	Mmu.17556	XR_092351	LOC100426 120	BCL2/adenovirus E1B 19 kDa protein-interacting protein 3-like
E03	N/A	XM 002803495	LOC693554	Serine/threonine-protein kinase B-raf-like
E04	N/A	XM 001083217	LOC695164	Tumor protein p73-like
E05	Mmu.8758	XM 002802530	LOC698160	Tumor necrosis factor receptor type 1-associated DEATH domain protein-like
E06	N/A	XM 002804163	LOC698788	Caspase-6-like
E07	N/A	XM 001095818	LOC707407	Apoptosis regulator Bcl-2-like
E08	N/A	XM 001100468	LOC708606	Protein FADD-like
E09	Mmu.13649	XM 001104846	LOC714869	B-cell lymphoma/leukemia 10-like
E10	N/A	XM 001107490	LOC716650	Tumor necrosis factor receptor superfamily member 10B-like
E11	Mmu.10857	NM 001047148	LTA	Lymphotoxin alpha (TNF superfamily, member 1)
E12	Mmu.11472	XM 001103746	LTBR	Lymphotoxin beta receptor (TNFR superfamily, member 3)
F01	Mmu.13662	XM 001089600	MAPK1	Mitogen-activated protein kinase 1
F02	Mmu.4052	XM 001102283	MCL1	Myeloid cell leukemia sequence 1 (BCL2-related)
F03	N/A	NM 001142873	MYC	V-myc myelocytomatosis viral oncogene homolog (avian)
F04	N/A	XM 001093532	NAIP	NLR family, apoptosis inhibitory protein
F05	Mmu.10897	XM 001085719	NOD1	Nucleotide-binding oligomerization domain containing 1
F06	Mmu.14465	XM 001087572	NOL3	Nucleolar protein 3 (apoptosis repressor with CARD domain)
F07	N/A	XM 002800569	PRKCA	Protein kinase C, alpha
F08	Mmu.18725	XM_001112804	PYCARD	PYD and CARD domain containing
F09	Mmu.16218	XM_001112004 XM_001084687	RIPK2	Receptor-interacting serine-threonine kinase 2
F10	Mmu.3364	NM 001047149	TNF	Tumor necrosis factor
F11	N/A	XM 001107790	TNFRSF10A	Tumor necrosis factor receptor superfamily, member 10a
F12	N/A	XM_001107770	TNFRSF11B	Tumor necrosis factor receptor superfamily, member 11b
G01	Mmu.12049	XM_001070713	TNFRSF1A	Tumor necrosis factor receptor superfamily, member 11A
G02	Mmu.13423	XM_001110232	TNFRSF21	Tumor necrosis factor receptor superfamily, member 21
G02	Mmu.15187	XM_001103782 XM_001093543	TNFRSF25	Tumor necrosis factor receptor superfamily, member 25
G03	N/A	XM_001093343	TNFRSF9	Tumor necrosis factor receptor superfamily, member 9
G05	Mmu.1012	XM_001090100	TNFSF10	Tumor necrosis factor (ligand) superfamily, member 10
G05	N/A	XM_001084788 XM_001099524	TNFSF8	Tumor necrosis factor (ligand) superfamily, member 10 Tumor necrosis factor (ligand) superfamily, member 8
G07	Mmu.3286	NM 001047151	TP53	Tumor protein p53
G08	Mmu.12448	XM 001093747	TP53BP2	Tumor protein p53 binding protein, 2
G09	Mmu.1370	XM_001093747 XM_001082535	TRAF3	TNF receptor-associated factor 3
G10	N/A	XM_001082535 XM_001107048	TRAF4	TNF receptor-associated factor 3 TNF receptor-associated factor 4
G10 G11	Mmu.15706	XM_001107048	TRAF5	TNF receptor-associated factor 4 TNF receptor-associated factor 5
G12	Mmu.15/06 N/A	XM_001108970 XM_001114854	XIAP	X-linked inhibitor of apoptosis
H01	Mmu.4974	NM 001033084	ACTB	A-linked inhibitor of apoprosis Actin, beta
H02	Mmu.5037	NM 001033084	B2M	,
H02	Mmu.3145	XM 001105471	GAPDH	Beta-2-microglobulin
H03	Mmu.3145 Mmu.12316	XM_001105471 XM_001097691	LOC709186	Glyceraldehyde-3-phosphate dehydrogenase
H04 H05	Mmu.12316 Mmu.2512	XM_001097691 XM_001115079	RPL13A	Hypoxanthine-guanine phosphoribosyltransferase-like
H05		_	QGDC	Ribosomal protein L13A
	N/A	SA_00125	-	Rhesus Macaque Genomic DNA Contamination
H07	N/A	SA_00104	RTC	Reverse Transcription Control
H08	N/A	SA_00104	RTC	Reverse Transcription Control
H09	N/A	SA_00104	RTC	Reverse Transcription Control
H10	N/A	SA_00103	PPC	Positive PCR Control
H11	N/A	SA_00103	PPC	Positive PCR Control
H12	N/A	SA_00103	PPC	Positive PCR Control

Related products

For optimal performance, RT² Profiler PCR Arrays should be used together with the RT² First Strand Kit for cDNA synthesis and RT2 SYBR[®] Green qPCR Mastermixes for PCR.

Product	Contents	Cat. no.
RT ² First Strand Kit (12)	Enzymes and reagents for cDNA synthesis	330401
RT ² SYBR Green qPCR Mastermix (2)*	For 2 x 96 assays in 96-well plates; suitable for use with real-time cyclers that do not require a reference dye, including: Bio-Rad models CFX96, CFX384, DNA Engine Opticon 2; Bio-Rad/MJ Research Chromo4; Roche LightCycler 480 (96-well and 384-well); all other cyclers	330500
RT ² SYBR Green ROX [™] qPCR Mastermix (2)*	For 2 x 96 assays in 96-well plates; suitable for use with the following real-time cyclers: Applied Biosystems models 5700, 7000, 7300, 7500 [Standard and FAST], 7700, 7900HT 96-well block [Standard and FAST] and 384-well block, StepOnePlus; Eppendorf Mastercycler ep realplex models 2, 2S, 4, 4S; Stratagene models Mx3000P, Mx3005P, Mx4000; Takara TP-800	330520
RT ² SYBR Green Fluor qPCR Mastermix (2)*	For 2 x 96 assays in 96-well plates; suitable for use with the following real-time cyclers: Bio-Rad models iCycler, iQ5, MyiQ, MyiQ2	330510

^{*} Larger kit sizes available; please inquire.

RT² Profiler PCR Array products are intended for molecular biology applications. These products are not intended for the diagnosis, prevention, or treatment of a disease.

For up-to-date licensing information and product-specific disclaimers, see the respective QIAGEN kit handbook or user manual. QIAGEN kit handbooks and user manuals are available at www.qiagen. com or can be requested from QIAGEN Technical Services or your local distributor.

Trademarks: QIAGEN® (QIAGEN Group); Applied Biosystems®, ViiA™, StepOnePlus™, ROX™ (Applera Corporation or its subsidiaries); Bio-Rad®, iCycler®, iQ™, MyiQ™, Chromo4™, CFX96™, DNA Engine Opticon®, CFX384™ (Bio-Rad Laboratories, Inc.)Stratagene®, Mx3005P®, Mx3000P®, Mx4000® (Stratagene); Eppendorf®, Mastercycler® (Eppendorf AG); Roche®, LightCycler® (Roche Group); Fluidigm[®] BioMark[™] (Fluidigm Corporation); SYBR[®] (Molecular Probes, Inc.). 1066029 03/2011 © 2011 QIAGEN, all rights reserved.

Canada • 800-572-9613 www.aiaaen.com China • 8621-3865-3865 Denmark ■ 80-885945 Australia • 1-800-243-800 Finland • 0800-914416 Austria • 0800/281010 France • 01-60-920-930 Belgium • 0800-79612 Germany ■ 02103-29-12000 Brazil • 0800-557779 Hong Kong • 800 933 965

Ireland = 1800 555 049 Italy • 800-787980 Japan • 03-6890-7300 Korea (South) • 080-000-7145 Luxembourg ■ 8002 2076 Mexico = 01-800-7742-436 The Netherlands • 0800 0229592 USA • 800-426-8157

Norway ■ 800-18859 Singapore ■ 1800-742-4368 Spain ■ 91-630-7050 Sweden • 020-790282 Switzerland • 055-254-22-11 UK • 01293-422-911

