

RT² Profiler PCR Array (96-Well Format and 384-Well [4 x 96] Format)

Rat T-Cell & B-Cell Activation

Cat. no. 330231 PARN-053ZA

For pathway expression analysis

Format	For use with the following real-time cyclers
RT ² Profiler PCR Array, Format A	Applied Biosystems® models 5700, 7000, 7300, 7500, 7700, 7900HT, ViiA™ 7 (96-well block); Bio-Rad® models iCycler®, iQ™ 5, MyiQ™, MyiQ2; Bio-Rad/MJ Research Chromo4™; Eppendorf® Mastercycler® ep realplex models 2, 2s, 4, 4s; Stratagene® models Mx3005P®, Mx3000P®; Takara TP-800
RT ² Profiler PCR Array, Format C	Applied Biosystems models 7500 (Fast block), 7900HT (Fast block), StepOnePlus™, ViiA 7 (Fast block)
RT ² Profiler PCR Array, Format D	Bio-Rad CFX96™; Bio-Rad/MJ Research models DNA Engine Opticon®, DNA Engine Opticon 2; Stratagene Mx4000®
RT ² Profiler PCR Array, Format E	Applied Biosystems models 7900HT (384-well block), ViiA 7 (384-well block); Bio-Rad CFX384™
RT ² Profiler PCR Array, Format F	Roche® LightCycler® 480 (96-well block)
RT ² Profiler PCR Array, Format G	Roche LightCycler 480 (384-well block)
RT ² Profiler PCR Array, Format H	Fluidigm® BioMark™



Sample & Assay Technologies

Description

The Rat T-Cell & B-Cell Activation RT² Profiler PCR Array profiles the expression of 84 genes representing T Cell and B Cell Activation, a key part of adaptive immunity. This array includes genes involved B-cell activation as well as genes involved in B-cell proliferation and differentiation. Genes involved in the activation of T-cells and their proliferation and differentiation are also represented along with genes regulating Th1 and Th2 development and T-cell polarization. Genes involved in the activation of macrophages, neutrophils, and natural killer cells are also included. Using real-time PCR, you can easily and reliably analyze expression of a focused panel of genes related to T Cell and B Cell Activation with this array.

For further details, consult the *RT² Profiler PCR Array Handbook*.

Shipping and storage

RT² Profiler PCR Arrays in formats A, C, D, E, F, and G are shipped at ambient temperature, on dry ice, or blue ice packs depending on destination and accompanying products. RT² Profiler PCR Arrays in format H are shipped on dry ice or blue ice packs.

For long term storage, keep plates at –20°C.

Note: Ensure that you have the correct RT² Profiler PCR Array format for your real-time cycler (see table above).

Note: Open the package and store the products appropriately immediately on receipt.

Array layout (96-well)

For 384-well 4 x 96 PCR arrays, genes are present in a staggered format. Refer to the *RT² Profiler PCR Array Handbook* for layout.

	1	2	3	4	5	6	7	8	9	10	11	12
A	Ada	Apc	Bad	Bcl2	Casp3	Cblb	Cd3	Cd4	Ccnd3	Ccr3	Ccr4	Cd1d1
B	Cd2	Cd276	Cd28	Cd38	Cd3d	Cd3e	Cd3g	Cd4	Cd40	Cd40lg	Cd47	Cd48
C	Cd74	Cd79a	Cd81	Cd8a	Cd8b	Cdkn1a	Cr2	Csf2	Ctla4	Cxcl12	Cxcr4	Cxcr5
D	Dpp4	Egr1	Flt3	Foxp3	Hsp90aa1	Icam1	Ifng	Il10	Il11	Il12b	Il15	Il18
E	Il18r1	Il1b	Il2	Il27	Il2ra	Il4	Il6	Il7	Inha	Irf4	Jag2	Lck
F	Map3k7	Ms4a1	Nck1	Nos2	Nr3c1	Pax1	Pdcd1lg2	Pik3r1	Prkcd	Prkcx	Prlr	Ptprc
G	Rag1	Socs1	Spp1	Tgfb1	Tlr1	Tlr4	Tlr6	Tnfrsf14	Tp53	Traf6	Vav1	Was
H	Actb	B2m	Hprt1	Ldha	Rplp1	RGDC	RTC	RTC	RTC	PPC	PPC	PPC

Gene table: RT² Profiler PCR Array

Position	UniGene	GenBank	Symbol	Description
A01	Rn.12689	NM_130399	Ada	Adenosine deaminase
A02	Rn.88057	NM_012499	Apc	Adenomatous polyposis coli
A03	Rn.36696	NM_022698	Bad	BCL2-associated agonist of cell death
A04	Rn.9996	NM_016993	Bcl2	B-cell CLL/lymphoma 2
A05	Rn.10562	NM_012922	Casp3	Caspase 3
A06	Rn.21799	NM_133601	Cblb	Cas-Br-M (murine) ecotropic retroviral transforming sequence b
A07	Rn.10139	NM_013025	Cd3	Chemokine (C-C motif) ligand 3
A08	Rn.37880	NM_053858	Cd4	Chemokine (C-C motif) ligand 4
A09	Rn.3483	NM_012766	Ccnd3	Cyclin D3
A10	Rn.9890	NM_053958	Ccr3	Chemokine (C-C motif) receptor 3
A11	Rn.81076	NM_133532	Ccr4	Chemokine (C-C motif) receptor 4
A12	Rn.11120	NM_017079	Cd1d1	CD1d1 molecule
B01	Rn.10328	NM_012830	Cd2	Cd2 molecule
B02	Rn.76129	NM_182824	Cd276	Cd276 molecule
B03	Rn.10327	NM_013121	Cd28	Cd28 molecule
B04	Rn.11414	NM_013127	Cd38	CD38 molecule
B05	Rn.10329	NM_013169	Cd3d	CD3 molecule, delta
B06	Rn.96221	NM_001108140	Cd3e	CD3 molecule, epsilon
B07	Rn.202653	NM_001077646	Cd3g	CD3 molecule, gamma
B08	Rn.10748	NM_012705	Cd4	Cd4 molecule
B09	Rn.25180	NM_134360	Cd40	CD40 molecule, TNF receptor superfamily member 5
B10	Rn.44218	NM_053353	Cd40lg	CD40 ligand
B11	Rn.7409	NM_019195	Cd47	Cd47 molecule
B12	Rn.3705	NM_139103	Cd48	Cd48 molecule
C01	Rn.33804	NM_013069	Cd74	Cd74 molecule, major histocompatibility complex, class II invariant chain
C02	Rn.178258	XM_001060872	Cd79a	Cd79a molecule, immunoglobulin-associated alpha
C03	Rn.1975	NM_013087	Cd81	Cd81 molecule
C04	Rn.10306	NM_031538	Cd8a	CD8a molecule
C05	Rn.10330	NM_031539	Cd8b	CD8b molecule
C06	Rn.10089	NM_080782	Cdkn1a	Cyclin-dependent kinase inhibitor 1A
C07	Rn.23490	NM_001105989	Cr2	Complement receptor 2
C08	Rn.44285	XM_340799	Csf2	Colony stimulating factor 2 (granulocyte-macrophage)
C09	Rn.10259	NM_031674	Ctla4	Cytotoxic T-lymphocyte-associated protein 4
C10	Rn.54439	NM_022177	Cxcl12	Chemokine (C-X-C motif) ligand 12 (stromal cell-derived factor 1)
C11	Rn.44431	NM_022205	Cxcr4	Chemokine (C-X-C motif) receptor 4
C12	Rn.127121	NM_053303	Cxcr5	Chemokine (C-X-C motif) receptor 5
D01	Rn.91364	NM_012789	Dpp4	Dipeptidylpeptidase 4
D02	Rn.9096	NM_012551	Egr1	Early growth response 1
D03	Rn.6774	NM_001100822	Flt3	Fms-related tyrosine kinase 3
D04	Rn.177272	NM_001108250	Foxp3	Forkhead box P3
D05	Rn.119867	NM_175761	Hsp90aa1	Heat shock protein 90, alpha (cytosolic), class A member 1
D06	Rn.12	NM_012967	Icam1	Intercellular adhesion molecule 1
D07	Rn.10795	NM_138880	Ifng	Interferon gamma
D08	Rn.9868	NM_012854	Il10	Interleukin 10
D09	Rn.198483	NM_133519	Il11	Interleukin 11

Position	UniGene	GenBank	Symbol	Description
D10	Rn.48686	NM_022611	Il12b	Interleukin 12b
D11	Rn.2490	NM_013129	Il15	Interleukin 15
D12	Rn.11118	NM_019165	Il18	Interleukin 18
E01	Rn.149181	NM_001106905	Il18r1	Interleukin 18 receptor 1
E02	Rn.9869	NM_031512	Il1b	Interleukin 1 beta
E03	Rn.9871	NM_053836	Il2	Interleukin 2
E04	N/A	XM_344962	Il27	Interleukin 27
E05	Rn.9872	NM_013163	Il2ra	Interleukin 2 receptor, alpha
E06	Rn.108255	NM_201270	Il4	Interleukin 4
E07	Rn.9873	NM_012589	Il6	Interleukin 6
E08	Rn.10793	NM_013110	Il7	Interleukin 7
E09	Rn.8831	NM_012590	Inha	Inhibin alpha
E10	Rn.82114	NM_001106108	Irf4	Interferon regulatory factor 4
E11	Rn.22459	XM_343119	Jag2	Jagged 2
E12	Rn.22791	NM_001100709	Lck	Lymphocyte-specific protein tyrosine kinase
F01	Rn.24019	NM_001107920	Map3k7	Mitogen activated protein kinase kinase kinase 7
F02	Rn.16385	NM_001107578	Ms4a1	Membrane-spanning 4-domains, subfamily A, member 1
F03	Rn.8768	NM_001106851	Nck1	NCK adaptor protein 1
F04	Rn.10400	NM_012611	Nos2	Nitric oxide synthase 2, inducible
F05	Rn.90070	NM_012576	Nr3c1	Nuclear receptor subfamily 3, group C, member 1
F06	Rn.40296	NM_001107787	Pax1	Paired box 1
F07	Rn.130112	NM_001107582	Pdcd1lg2	Programmed cell death 1 ligand 2
F08	Rn.10599	NM_013005	Pik3r1	Phosphoinositide-3-kinase, regulatory subunit 1 (alpha)
F09	Rn.98279	NM_133307	Prkcd	Protein kinase C, delta
F10	Rn.8227	XM_341553	Prkcg	Protein kinase C, theta
F11	Rn.9757	NM_012630	Prlr	Prolactin receptor
F12	Rn.90166	NM_001109887	Ptrc	Protein tyrosine phosphatase, receptor type, C
G01	Rn.92344	NM_053468	Rag1	Recombination activating gene 1
G02	Rn.82754	NM_145879	Socs1	Suppressor of cytokine signaling 1
G03	Rn.8871	NM_012881	Spp1	Secreted phosphoprotein 1
G04	Rn.40136	NM_021578	Tgfb1	Transforming growth factor, beta 1
G05	Rn.107212	NM_001172120	Tlr1	Toll-like receptor 1
G06	Rn.14534	NM_019178	Tlr4	Toll-like receptor 4
G07	Rn.163249	NM_207604	Tlr6	Toll-like receptor 6
G08	N/A	XM_236794	Tnfsf14	Tumor necrosis factor (ligand) superfamily, member 14
G09	Rn.54443	NM_030989	Tp53	Tumor protein p53
G10	Rn.220435	NM_001107754	Traf6	Tnf receptor-associated factor 6
G11	Rn.48861	NM_012759	Vav1	Vav 1 guanine nucleotide exchange factor
G12	Rn.207069	NM_001108248	Was	Wiskott-Aldrich syndrome homolog (human)
H01	Rn.94978	NM_031144	Actb	Actin, beta
H02	Rn.1868	NM_012512	B2m	Beta-2 microglobulin
H03	Rn.47	NM_012583	Hprt1	Hypoxanthine phosphoribosyltransferase 1
H04	Rn.107896	NM_017025	Ldha	Lactate dehydrogenase A
H05	Rn.973	NM_001007604	Rplp1	Ribosomal protein, large, P1
H06	N/A	U26919	RGDC	Rat Genomic DNA Contamination
H07	N/A	SA_00104	RTC	Reverse Transcription Control
H08	N/A	SA_00104	RTC	Reverse Transcription Control
H09	N/A	SA_00104	RTC	Reverse Transcription Control
H10	N/A	SA_00103	PPC	Positive PCR Control
H11	N/A	SA_00103	PPC	Positive PCR Control
H12	N/A	SA_00103	PPC	Positive PCR Control

Related products

For optimal performance, RT² Profiler PCR Arrays should be used together with the RT² First Strand Kit for cDNA synthesis and RT² SYBR[®] Green qPCR Mastermixes for PCR.

Product	Contents	Cat. no.
RT ² First Strand Kit (12)	Enzymes and reagents for cDNA synthesis	330401
RT ² SYBR Green qPCR Mastermix (2)*	For 2 x 96 assays in 96-well plates; suitable for use with real-time cyclers that do not require a reference dye, including: Bio-Rad models CFX96, CFX384, DNA Engine Opticon 2; Bio-Rad/MJ Research Chromo4; Roche LightCycler 480 (96-well and 384-well); all other cyclers	330500
RT ² SYBR Green ROX [™] qPCR Mastermix (2)*	For 2 x 96 assays in 96-well plates; suitable for use with the following real-time cyclers: Applied Biosystems models 5700, 7000, 7300, 7500 [Standard and FAST], 7700, 7900HT 96-well block [Standard and FAST] and 384-well block, StepOnePlus; Eppendorf Mastercycler ep realplex models 2, 2S, 4, 4S; Stratagene models Mx3000P, Mx3005P, Mx4000; Takara TP-800	330520
RT ² SYBR Green Fluor qPCR Mastermix (2)*	For 2 x 96 assays in 96-well plates; suitable for use with the following real-time cyclers: Bio-Rad models iCycler, iQ5, MyiQ, MyiQ2	330510

* Larger kit sizes available; please inquire.

RT² Profiler PCR Array products are intended for molecular biology applications. These products are not intended for the diagnosis, prevention, or treatment of a disease.

For up-to-date licensing information and product-specific disclaimers, see the respective QIAGEN kit handbook or user manual. QIAGEN kit handbooks and user manuals are available at www.qiagen.com or can be requested from QIAGEN Technical Services or your local distributor.

Trademarks: QIAGEN® (QIAGEN Group); Applied Biosystems®, ViiA™, StepOnePlus™, ROX™ (Applied Biosystems Corporation or its subsidiaries); Bio-Rad®, iCycler®, iQ™, MyiQ™, Chromo4™, CFX96™, DNA Engine Opticon®, CFX384™ (Bio-Rad Laboratories, Inc.); Stratagene®, Mx3005P®, Mx3000P®, Mx4000® (Stratagene); Eppendorf®, Mastercycler® (Eppendorf AG); Roche®, LightCycler® (Roche Group); Fluidigm® BioMark™ (Fluidigm Corporation); SYBR® (Molecular Probes, Inc.).

1066029 03/2011 © 2011 QIAGEN, all rights reserved.

www.qiagen.com

Canada ■ 800-572-9613

China ■ 8621-3865-3865

Denmark ■ 80-885945

Finland ■ 0800-914416

France ■ 01-60-920-930

Germany ■ 02103-29-12000

Hong Kong ■ 800 933 965

Ireland ■ 1800 555 049

Italy ■ 800-787980

Japan ■ 03-6890-7300

Korea (South) ■ 080-000-7145

Luxembourg ■ 8002 2076

Mexico ■ 01-800-7742-436

The Netherlands ■ 0800 0229592

Norway ■ 800-18859

Singapore ■ 1800-742-4368

Spain ■ 91-630-7050

Sweden ■ 020-790282

Switzerland ■ 055-254-22-11

UK ■ 01293-422-911

USA ■ 800-426-8157

Australia ■ 1-800-243-800

Austria ■ 0800/281010

Belgium ■ 0800-79612

Brazil ■ 0800-557779



Sample & Assay Technologies