

RT² Profiler PCR Array (Rotor-Gene[®] Format)

Rat Fibrosis

Cat. no. 330231 PARN-120ZR

For pathway expression analysis

Format	For use with the following real-time cyclers
RT ² Profiler PCR Array, Format R	Rotor-Gene Q, other Rotor-Gene cyclers

Description

The Rat Fibrosis RT² Profiler PCR Array profiles the expression of 84 key genes involved in dysregulated tissue remodeling during the repair and healing of wounds. Wound healing consists of three phases: inflammation, granulation and tissue remodeling. During tissue remodeling, apoptosis removes myofibroblasts, and the deposited extracellular matrix (ECM) is remodeled to resemble the original tissue. Fibrosis occurs when inappropriate tissue remodeling results in excess ECM deposition due to inappropriate survival of myofibroblasts or lack of ECM proteolytic degradation. The inflammation and enhanced TGF β signaling often present in fibrotic tissues causes cells to differentiate into myofibroblasts via epithelial-to-mesenchymal transition (EMT). On the other side of the spectrum, chronic wounds feature dysregulated tissue remodeling with enhanced ECM degradation. This array contains genes encoding ECM remodeling enzymes, TGF β signaling molecules and inflammatory cytokines, as well as additional genes important for fibrosis. Using real-time PCR, you can easily and reliably analyze the expression of a focused panel of genes involved in fibrosis and chronic wounds with this array.

For further details, consult the *RT² Profiler PCR Array Handbook*.

Shipping and storage

RT² Profiler PCR Arrays in the Rotor-Gene format are shipped at ambient temperature, on dry ice, or blue ice packs depending on destination and accompanying products.

For long term storage, keep plates at -20°C .

Note: Ensure that you have the correct RT² Profiler PCR Array format for your real-time cycler (see table above).

Note: Open the package and store the products appropriately immediately on receipt.



Sample & Assay Technologies

Array layout

The 96 real-time assays in the Rotor-Gene format are located in wells 1–96 of the Rotor-Disc™ (plate A1–A12=Rotor-Disc 1–12, plate B1–B12=Rotor-Disc 13–24, etc.). To maintain data analysis compatibility, wells 97–100 do not contain real-time assays but will contain master mix to account for weight balance.

Gene table: RT² Profiler PCR Array

Position	UniGene	GenBank	Symbol	Description
A01	Rn.195319	NM_031004	Acta2	Smooth muscle alpha-actin
A02	Rn.6319	NM_134432	Agt	Angiotensinogen (serpin peptidase inhibitor, clade A, member 8)
A03	Rn.11422	NM_033230	Akt1	V-akt murine thymoma viral oncogene homolog 1
A04	Rn.9996	NM_016993	Bcl2	B-cell CLL/lymphoma 2
A05	Rn.18030	XM_342591	Bmp7	Bone morphogenetic protein 7
A06	Rn.22518	NM_031556	Cav1	Caveolin 1, caveolae protein
A07	Rn.10632	NM_019205	Ccl11	Chemokine (C-C motif) ligand 11
A08	Rn.137780	NM_001105822	Ccl12	Chemokine (C-C motif) ligand 12
A09	Rn.10139	NM_013025	Ccl3	Chemokine (C-C motif) ligand 3
A10	Rn.211983	NM_021866	Ccr2	Chemokine (C-C motif) receptor 2
A11	Rn.6479	NM_024125	Cebpb	CCAAT/enhancer binding protein (C/EBP), beta
A12	Rn.107239	NM_053356	Col1a2	Collagen, type I, alpha 2
B01	Rn.3247	NM_032085	Col3a1	Collagen, type III, alpha 1
B02	Rn.17145	NM_022266	Ctgf	Connective tissue growth factor
B03	Rn.44431	NM_022205	Cxcr4	Chemokine (C-X-C motif) receptor 4
B04	Rn.106103	NM_024129	Dcn	Decorin
B05	Rn.10918	NM_012548	Edn1	Endothelin 1
B06	Rn.6075	NM_012842	Egf	Epidermal growth factor
B07	Rn.187025	NM_001010968	Eng	Endoglin
B08	Rn.9725	NM_012908	Faslg	Fas ligand (TNF superfamily, member 6)
B09	Rn.42929	NM_019282	Grem1	Gremlin 1, cysteine knot superfamily, homolog (Xenopus laevis)
B10	Rn.10468	NM_017017	Hgf	Hepatocyte growth factor
B11	Rn.10795	NM_138880	Irfng	Interferon gamma
B12	Rn.9868	NM_012854	Il10	Interleukin 10
C01	Rn.9921	NM_053828	Il13	Interleukin 13
C02	Rn.162308	NM_133538	Il13ra2	Interleukin 13 receptor, alpha 2
C03	Rn.12300	NM_017019	Il1a	Interleukin 1 alpha
C04	Rn.9869	NM_031512	Il1b	Interleukin 1 beta
C05	Rn.108255	NM_201270	Il4	Interleukin 4
C06	Rn.44227	NM_021834	Il5	Interleukin 5
C07	Rn.95042	NM_133409	Ilk	Integrin-linked kinase
C08	Rn.30020	NM_031815	Inhbe	Inhibin beta E
C09	Rn.91044	NM_030994	Itga1	Integrin, alpha 1
C10	Rn.83597	XM_345156	Itga2	Integrin, alpha 2
C11	Rn.154664	NM_001108292	Itga3	Integrin, alpha 3
C12	Rn.23339	NM_001106549	Itgav	Integrin, alpha V
D01	Rn.25733	NM_017022	Itgb1	Integrin, beta 1
D02	Rn.162202	NM_153720	Itgb3	Integrin, beta 3
D03	Rn.16988	NM_147139	Itgb5	Integrin, beta 5
D04	Rn.19828	NM_001004263	Itgb6	Integrin, beta 6
D05	Rn.212213	NM_001108726	Itgb8	Integrin, beta 8
D06	Rn.93714	NM_021835	Jun	Jun oncogene
D07	Rn.11372	NM_017061	Lox	Lysyl oxidase
D08	Rn.40942	NM_021587	Libp1	Latent transforming growth factor beta binding protein 1
D09	Rn.10997	NM_133530	Mmp13	Matrix metalloproteinase 13
D10	Rn.10371	NM_031056	Mmp14	Matrix metalloproteinase 14 (membrane-inserted)
D11	Rn.79007	NM_001134530	Mmp1a	Matrix metalloproteinase 1a (interstitial collagenase)
D12	Rn.6422	NM_031054	Mmp2	Matrix metalloproteinase 2
E01	Rn.32086	NM_133523	Mmp3	Matrix metalloproteinase 3
E02	Rn.44474	NM_022221	Mmp8	Matrix metalloproteinase 8
E03	Rn.10209	NM_031055	Mmp9	Matrix metalloproteinase 9
E04	Rn.12072	NM_012603	Myc	Myelocytomatosis oncogene
E05	Rn.2411	XM_342346	Nfkb1	Nuclear factor of kappa light polypeptide gene enhancer in B-cells 1
E06	Rn.10999	NM_012801	Pdgfa	Platelet-derived growth factor alpha polypeptide
E07	Rn.198230	NM_031524	Pdgfb	Platelet-derived growth factor beta polypeptide (simian sarcoma viral (v-sis) oncogene homolog)
E08	Rn.107102	NM_013151	Plat	Plasminogen activator, tissue

Position	UniGene	GenBank	Symbol	Description
E09	Rn.6064	NM_013085	Plau	Plasminogen activator, urokinase
E10	Rn.20178	NM_053491	Plg	Plasminogen
E11	Rn.1419	NM_022519	Serpina1	Serpin peptidase inhibitor, clade A (alpha-1 antiproteinase, antitrypsin), member 1
E12	Rn.29367	NM_012620	Serpine1	Serpin peptidase inhibitor, clade E (nexin, plasminogen activator inhibitor type 1), member 1
F01	Rn.98199	NM_017173	Serpinh1	Serine (or cysteine) peptidase inhibitor, clade H, member 1
F02	Rn.2755	NM_019191	Smad2	SMAD family member 2
F03	Rn.10636	NM_013095	Smad3	SMAD family member 3
F04	Rn.9774	NM_019275	Smad4	SMAD family member 4
F05	Rn.42472	NM_001109002	Smad6	SMAD family member 6
F06	Rn.29980	NM_030858	Smad7	SMAD family member 7
F07	Rn.8008	NM_053805	Snai1	Snail homolog 1 (Drosophila)
F08	Rn.44609	NM_012655	Sp1	Sp1 transcription factor
F09	Rn.33229	NM_032612	Stat1	Signal transducer and activator of transcription 1
F10	Rn.6880	NM_001044250	Stat6	Signal transducer and activator of transcription 6
F11	Rn.40136	NM_021578	Tgfb1	Transforming growth factor, beta 1
F12	Rn.24539	NM_031131	Tgfb2	Transforming growth factor, beta 2
G01	Rn.7018	NM_013174	Tgfb3	Transforming growth factor, beta 3
G02	Rn.44402	NM_012775	Tgfb1	Transforming growth factor, beta receptor 1
G03	Rn.9954	NM_031132	Tgfb2	Transforming growth factor, beta receptor II
G04	Rn.154349	NM_001015020	Tgif1	TGFB-induced factor homeobox 1
G05	Rn.185771	NM_001013062	Thbs1	Thrombospondin 1
G06	Rn.165619	NM_001169138	Thbs2	Thrombospondin 2
G07	Rn.25754	NM_053819	Timp1	TIMP metalloproteinase inhibitor 1
G08	Rn.10161	NM_021989	Timp2	TIMP metalloproteinase inhibitor 2
G09	Rn.119634	NM_012886	Timp3	TIMP metalloproteinase inhibitor 3
G10	Rn.155651	NM_001109393	Timp4	Tissue inhibitor of metalloproteinase 4
G11	Rn.2275	NM_012675	Tnf	Tumor necrosis factor (TNF superfamily, member 2)
G12	Rn.1923	NM_031836	Vegfa	Vascular endothelial growth factor A
H01	Rn.94978	NM_031144	Actb	Actin, beta
H02	Rn.1868	NM_012512	B2m	Beta-2 microglobulin
H03	Rn.47	NM_012583	Hprt1	Hypoxanthine phosphoribosyltransferase 1
H04	Rn.107896	NM_017025	Ldha	Lactate dehydrogenase A
H05	Rn.973	NM_001007604	Rplp1	Ribosomal protein, large, P1
H06	N/A	U26919	RGDC	Rat Genomic DNA Contamination
H07	N/A	SA_00104	RTC	Reverse Transcription Control
H08	N/A	SA_00104	RTC	Reverse Transcription Control
H09	N/A	SA_00104	RTC	Reverse Transcription Control
H10	N/A	SA_00103	PPC	Positive PCR Control
H11	N/A	SA_00103	PPC	Positive PCR Control
H12	N/A	SA_00103	PPC	Positive PCR Control

Related products

For optimal performance, RT² Profiler PCR Arrays should be used together with the RT² First Strand Kit for cDNA synthesis and RT² SYBR[®] Green qPCR Mastermixes for PCR.

Product	Contents	Cat. no.
RT ² First Strand Kit (12)	Enzymes and reagents for cDNA synthesis	330401
RT ² SYBR Green ROX™ FAST Mastermix (2)*	For 2 x 96 assays in 96-well plates; suitable for use with the Rotor-Gene Q and other Rotor-Gene cyclers	330620

* Larger kit sizes available; please inquire.

RT² Profiler PCR Array products are intended for molecular biology applications. These products are not intended for the diagnosis, prevention, or treatment of a disease.

For up-to-date licensing information and product-specific disclaimers, see the respective QIAGEN kit handbook or user manual. QIAGEN kit handbooks and user manuals are available at www.qiagen.com or can be requested from QIAGEN Technical Services or your local distributor.

Trademarks: QIAGEN[®], Rotor-Gene[®], Rotor-Disc[™] (QIAGEN Group); ROX[™] (Applied Biosystems Corporation or its subsidiaries); SYBR[®] (Molecular Probes, Inc.).

1067688 03/2011 © 2011 QIAGEN, all rights reserved.

www.qiagen.com

Canada ■ 800-572-9613

China ■ 8621-3865-3865

Denmark ■ 80-885945

Finland ■ 0800-914416

France ■ 01-60-920-930

Germany ■ 02103-29-12000

Hong Kong ■ 800 933 965

Ireland ■ 1800 555 049

Italy ■ 800-787980

Japan ■ 03-6890-7300

Korea (South) ■ 080-000-7145

Luxembourg ■ 8002 2076

Mexico ■ 01-800-7742-436

The Netherlands ■ 0800 0229592

Norway ■ 800-18859

Singapore ■ 1800-742-4368

Spain ■ 91-630-7050

Sweden ■ 020-790282

Switzerland ■ 055-254-22-11

UK ■ 01293-422-911

USA ■ 800-426-8157

Australia ■ 1-800-243-800

Austria ■ 0800/281010

Belgium ■ 0800-79612

Brazil ■ 0800-557779



Sample & Assay Technologies