

RT² Profiler PCR Array (96-Well Format and 384-Well [4 x 96] Format)

Rat Wound Healing

Cat. no. 330231 PARN-121ZA

For pathway expression analysis

Format	For use with the following real-time cyclers
RT ² Profiler PCR Array, Format A	Applied Biosystems® models 5700, 7000, 7300, 7500, 7700, 7900HT, ViiA™ 7 (96-well block); Bio-Rad® models iCycler®, iQ™ 5, MyiQ™, MyiQ2; Bio-Rad/MJ Research Chromo4™; Eppendorf® Mastercycler® ep realplex models 2, 2s, 4, 4s; Stratagene® models Mx3005P®, Mx3000P®; Takara TP-800
RT ² Profiler PCR Array, Format C	Applied Biosystems models 7500 (Fast block), 7900HT (Fast block), StepOnePlus™, ViiA 7 (Fast block)
RT ² Profiler PCR Array, Format D	Bio-Rad CFX96™; Bio-Rad/MJ Research models DNA Engine Opticon®, DNA Engine Opticon 2; Stratagene Mx4000®
RT ² Profiler PCR Array, Format E	Applied Biosystems models 7900HT (384-well block), ViiA 7 (384-well block); Bio-Rad CFX384™
RT ² Profiler PCR Array, Format F	Roche® LightCycler® 480 (96-well block)
RT ² Profiler PCR Array, Format G	Roche LightCycler 480 (384-well block)
RT ² Profiler PCR Array, Format H	Fluidigm® BioMark™



Sample & Assay Technologies

Description

The Rat Wound Healing RT² Profiler PCR Array profiles the expression of 84 key genes central to the wound healing response. Wound healing progresses via three overlapping phases: inflammation, granulation and tissue remodeling. After cutaneous injury, a blood clot forms, and inflammatory cells infiltrate the wound, secreting cytokines and growth factors to promote the inflammation phase. During the granulation phase, fibroblasts and other cells differentiate into myofibroblasts, which deposit extracellular matrix (ECM) proteins. Simultaneously, angiogenesis occurs, and keratinocytes proliferate and migrate to close the wound. In the final tissue remodeling phase, apoptosis eliminates myofibroblasts and extraneous blood vessels, and the ECM is remodeled to resemble the original tissue. Dysregulation of this last tissue remodeling phase leads to fibrosis. This array contains genes important for each of the three phases of wound healing, including ECM remodeling factors, inflammatory cytokines and chemokines, as well as growth factors and major signaling molecules. Using real-time PCR, you can easily and reliably analyze the expression of a focused panel of genes involved in wound healing, tissue injury and repair with this array.

For further details, consult the *RT² Profiler PCR Array Handbook*.

Shipping and storage

RT² Profiler PCR Arrays in formats A, C, D, E, F, and G are shipped at ambient temperature, on dry ice, or blue ice packs depending on destination and accompanying products. RT² Profiler PCR Arrays in format H are shipped on dry ice or blue ice packs.

For long term storage, keep plates at –20°C.

Note: Ensure that you have the correct RT² Profiler PCR Array format for your real-time cycler (see table above).

Note: Open the package and store the products appropriately immediately on receipt.

Array layout (96-well)

For 384-well 4 x 96 PCR arrays, genes are present in a staggered format. Refer to the *RT² Profiler PCR Array Handbook* for layout.

	1	2	3	4	5	6	7	8	9	10	11	12
A	Acta2	Actc1	Angpt1	Ccl12	Ccl7	Cd40lg	Cdh1	Col14a1	Col1a1	Col1a2	Col3a1	Col4a1
B	Col4a3	Col5a1	Col5a2	Col5a3	Csf2	Csf3	Ctgf	Ctnnb1	Ctsq	Ctsk	Ctsl1	Cxcl1
C	Cxcl11	Cxcl3	Cxcl5	Egf	Egfr	F13a1	F3	Fga	Fgf10	Fgf2	Fgf7	Hbegf
D	Hgf	Ifng	Igf1	Il10	Il1b	Il2	Il4	Il6	Il6st	Ilga1	Ilga2	Ilga3
E	Ilga4	Ilga5	Ilga6	Ilgav	Ilgb1	Ilgb3	Ilgb5	Ilgb6	Mapk1	Mapk3	Mif	Mmp1a
F	Mmp2	Mmp7	Mmp9	Pdgfra	Plat	Plau	Plaur	Plg	Pten	Ptgs2	Rac1	Rhoa
G	Serpina1	Stat3	Tagln	Tgfa	Tgfb1	Tgfb3	Timp1	Tnf	Vegfa	Vtn	Wisp1	Wnt5a
H	Actb	B2m	Hprt1	Ldha	Rplp1	RGDC	RTC	RTC	RTC	PPC	PPC	PPC

Gene table: RT² Profiler PCR Array

Position	UniGene	GenBank	Symbol	Description
A01	Rn.195319	NM_031004	Acta2	Smooth muscle alpha-actin
A02	Rn.3114	NM_019183	Actc1	Actin, alpha, cardiac muscle 1
A03	Rn.161953	NM_053546	Angpt1	Angiopietin 1
A04	Rn.137780	NM_001105822	Ccl12	Chemokine (C-C motif) ligand 12
A05	Rn.26815	NM_001007612	Ccl7	Chemokine (C-C motif) ligand 7
A06	Rn.44218	NM_053353	Cd40lg	CD40 ligand
A07	Rn.1303	NM_031334	Cdh1	Cadherin 1
A08	Rn.99441	NM_001130548	Col14a1	Collagen, type XIV, alpha 1
A09	Rn.2953	NM_053304	Col1a1	Collagen, type I, alpha 1
A10	Rn.107239	NM_053356	Col1a2	Collagen, type I, alpha 2
A11	Rn.3247	NM_032085	Col3a1	Collagen, type III, alpha 1
A12	Rn.53801	NM_001135009	Col4a1	Collagen, type IV, alpha 1
B01	Rn.121139	NM_001135759	Col4a3	Collagen, type IV, alpha 3
B02	Rn.117	NM_134452	Col5a1	Collagen, type V, alpha 1
B03	Rn.2875	NM_053488	Col5a2	Collagen, type V, alpha 2
B04	Rn.38654	NM_021760	Col5a3	Collagen, type V, alpha 3
B05	Rn.44285	XM_340799	Csf2	Colony stimulating factor 2 (granulocyte-macrophage)
B06	Rn.53973	NM_017104	Csf3	Colony stimulating factor 3 (granulocyte)
B07	Rn.17145	NM_022266	Ctgf	Connective tissue growth factor
B08	Rn.112601	NM_053357	Ctnnb1	Catenin (cadherin associated protein), beta 1
B09	Rn.103332	NM_001106041	Ctsq	Cathepsin G
B10	Rn.5598	NM_031560	Ctsk	Cathepsin K
B11	Rn.1294	NM_013156	Ctsl1	Cathepsin L1
B12	Rn.10907	NM_030845	Cxcl1	Chemokine (C-X-C motif) ligand 1 (melanoma growth stimulating activity, alpha)
C01	Rn.13664	NM_182952	Cxcl11	Chemokine (C-X-C motif) ligand 11
C02	Rn.10525	NM_138522	Cxcl3	Chemokine (C-X-C motif) ligand 3
C03	Rn.44449	NM_022214	Cxcl5	Chemokine (C-X-C motif) ligand 5
C04	Rn.6075	NM_012842	Egf	Epidermal growth factor
C05	Rn.37227	NM_031507	Egfr	Epidermal growth factor receptor
C06	Rn.42925	NM_021698	F13a1	Coagulation factor XIII, A1 polypeptide
C07	Rn.9980	NM_013057	F3	Coagulation factor III (thromboplastin, tissue factor)
C08	Rn.98846	NM_001008724	Fga	Fibrinogen alpha chain
C09	Rn.44439	NM_012951	Fgf10	Fibroblast growth factor 10
C10	Rn.31808	NM_019305	Fgf2	Fibroblast growth factor 2
C11	Rn.98842	NM_022182	Fgf7	Fibroblast growth factor 7
C12	Rn.10148	NM_012945	Hbegf	Heparin-binding EGF-like growth factor
D01	Rn.10468	NM_017017	Hgf	Hepatocyte growth factor
D02	Rn.10795	NM_138880	Ifng	Interferon gamma
D03	Rn.6282	NM_178866	Igf1	Insulin-like growth factor 1
D04	Rn.9868	NM_012854	Il10	Interleukin 10
D05	Rn.9869	NM_031512	Il1b	Interleukin 1 beta
D06	Rn.9871	NM_053836	Il2	Interleukin 2
D07	Rn.108255	NM_201270	Il4	Interleukin 4
D08	Rn.9873	NM_012589	Il6	Interleukin 6
D09	Rn.12138	NM_001008725	Il6st	Interleukin 6 signal transducer

Position	UniGene	GenBank	Symbol	Description
D10	Rn.91044	NM_030994	Itga1	Integrin, alpha 1
D11	Rn.83597	XM_345156	Itga2	Integrin, alpha 2
D12	Rn.154664	NM_001108292	Itga3	Integrin, alpha 3
E01	Rn.12704	NM_001107737	Itga4	Integrin, alpha 4
E02	Rn.100796	NM_001108118	Itga5	Integrin, alpha 5 (fibronectin receptor, alpha polypeptide)
E03	Rn.161799	XM_215984	Itga6	Integrin, alpha 6
E04	Rn.23339	NM_001106549	Itgav	Integrin, alpha V
E05	Rn.25733	NM_017022	Itgb1	Integrin, beta 1
E06	Rn.162202	NM_153720	Itgb3	Integrin, beta 3
E07	Rn.16988	NM_147139	Itgb5	Integrin, beta 5
E08	Rn.19828	NM_001004263	Itgb6	Integrin, beta 6
E09	Rn.34914	NM_053842	Mapk1	Mitogen activated protein kinase 1
E10	Rn.2592	NM_017347	Mapk3	Mitogen activated protein kinase 3
E11	Rn.2661	NM_031051	Mif	Macrophage migration inhibitory factor
E12	Rn.79007	NM_001134530	Mmp1a	Matrix metalloproteinase 1a (interstitial collagenase)
F01	Rn.6422	NM_031054	Mmp2	Matrix metalloproteinase 2
F02	Rn.10282	NM_012864	Mmp7	Matrix metalloproteinase 7
F03	Rn.10209	NM_031055	Mmp9	Matrix metalloproteinase 9
F04	Rn.10999	NM_012801	Pdgfa	Platelet-derived growth factor alpha polypeptide
F05	Rn.107102	NM_013151	Plat	Plasminogen activator, tissue
F06	Rn.6064	NM_013085	Plau	Plasminogen activator, urokinase
F07	Rn.82711	NM_017350	Plaur	Plasminogen activator, urokinase receptor
F08	Rn.20178	NM_053491	Plg	Plasminogen
F09	Rn.22158	NM_031606	Pten	Phosphatase and tensin homolog
F10	Rn.44369	NM_017232	Ptgs2	Prostaglandin-endoperoxide synthase 2
F11	Rn.29157	NM_134366	Rac1	Ras-related C3 botulinum toxin substrate 1
F12	Rn.107401	NM_057132	Rhoa	Ras homolog gene family, member A
G01	Rn.29367	NM_012620	Serpine1	Serpin peptidase inhibitor, clade E (nexin, plasminogen activator inhibitor type 1), member 1
G02	Rn.10247	NM_012747	Stat3	Signal transducer and activator of transcription 3
G03	Rn.34397	NM_031549	Tagln	Transgelin
G04	Rn.9952	NM_012671	Tgfa	Transforming growth factor alpha
G05	Rn.40136	NM_021578	Tgfb1	Transforming growth factor, beta 1
G06	Rn.9953	NM_017256	Tgfbr3	Transforming growth factor, beta receptor III
G07	Rn.25754	NM_053819	Timp1	TIMP metalloproteinase inhibitor 1
G08	Rn.2275	NM_012675	Tnf	Tumor necrosis factor (TNF superfamily, member 2)
G09	Rn.1923	NM_031836	Vegfa	Vascular endothelial growth factor A
G10	Rn.87493	NM_019156	Vtn	Vitronectin
G11	Rn.63486	NM_031716	Wisp1	WNT1 inducible signaling pathway protein 1
G12	Rn.48749	NM_022631	Wnt5a	Wingless-type MMTV integration site family, member 5A
H01	Rn.94978	NM_031144	Actb	Actin, beta
H02	Rn.1868	NM_012512	B2m	Beta-2 microglobulin
H03	Rn.47	NM_012583	Hprt1	Hypoxanthine phosphoribosyltransferase 1
H04	Rn.107896	NM_017025	Ldha	Lactate dehydrogenase A
H05	Rn.973	NM_001007604	Rplp1	Ribosomal protein, large, P1
H06	N/A	U26919	RGDC	Rat Genomic DNA Contamination
H07	N/A	SA_00104	RTC	Reverse Transcription Control
H08	N/A	SA_00104	RTC	Reverse Transcription Control
H09	N/A	SA_00104	RTC	Reverse Transcription Control
H10	N/A	SA_00103	PPC	Positive PCR Control
H11	N/A	SA_00103	PPC	Positive PCR Control
H12	N/A	SA_00103	PPC	Positive PCR Control

Related products

For optimal performance, RT² Profiler PCR Arrays should be used together with the RT² First Strand Kit for cDNA synthesis and RT² SYBR[®] Green qPCR Mastermixes for PCR.

Product	Contents	Cat. no.
RT ² First Strand Kit (12)	Enzymes and reagents for cDNA synthesis	330401
RT ² SYBR Green qPCR Mastermix (2)*	For 2 x 96 assays in 96-well plates; suitable for use with real-time cyclers that do not require a reference dye, including: Bio-Rad models CFX96, CFX384, DNA Engine Opticon 2; Bio-Rad/MJ Research Chromo4; Roche LightCycler 480 (96-well and 384-well); all other cyclers	330500
RT ² SYBR Green ROX™ qPCR Mastermix (2)*	For 2 x 96 assays in 96-well plates; suitable for use with the following real-time cyclers: Applied Biosystems models 5700, 7000, 7300, 7500 [Standard and FAST], 7700, 7900HT 96-well block [Standard and FAST] and 384-well block, StepOnePlus; Eppendorf Mastercycler ep realplex models 2, 2S, 4, 4S; Stratagene models Mx3000P, Mx3005P, Mx4000; Takara TP-800	330520
RT ² SYBR Green Fluor qPCR Mastermix (2)*	For 2 x 96 assays in 96-well plates; suitable for use with the following real-time cyclers: Bio-Rad models iCycler, iQ5, MyiQ, MyiQ2	330510

* Larger kit sizes available; please inquire.

RT² Profiler PCR Array products are intended for molecular biology applications. These products are not intended for the diagnosis, prevention, or treatment of a disease.

For up-to-date licensing information and product-specific disclaimers, see the respective QIAGEN kit handbook or user manual. QIAGEN kit handbooks and user manuals are available at www.qiagen.com or can be requested from QIAGEN Technical Services or your local distributor.

Trademarks: QIAGEN® (QIAGEN Group); Applied Biosystems®, ViiA™, StepOnePlus™, ROX™ (Applied Biosystems Corporation or its subsidiaries); Bio-Rad®, iCycler®, iQ™, MyiQ™, Chromo4™, CFX96™, DNA Engine Opticon®, CFX384™ (Bio-Rad Laboratories, Inc.); Stratagene®, Mx3005P®, Mx3000P®, Mx4000® (Stratagene); Eppendorf®, Mastercycler® (Eppendorf AG); Roche®, LightCycler® (Roche Group); Fluidigm® BioMark™ (Fluidigm Corporation); SYBR® (Molecular Probes, Inc.).

1066029 03/2011 © 2011 QIAGEN, all rights reserved.

www.qiagen.com

Canada ■ 800-572-9613

Ireland ■ 1800 555 049

Norway ■ 800-18859

China ■ 8621-3865-3865

Italy ■ 800-787980

Singapore ■ 1800-742-4368

Denmark ■ 80-885945

Japan ■ 03-6890-7300

Spain ■ 91-630-7050

Australia ■ 1-800-243-800

Finland ■ 0800-914416

Korea (South) ■ 080-000-7145

Sweden ■ 020-790282

Austria ■ 0800/281010

France ■ 01-60-920-930

Luxembourg ■ 8002 2076

Switzerland ■ 055-254-22-11

Belgium ■ 0800-79612

Germany ■ 02103-29-12000

Mexico ■ 01-800-7742-436

UK ■ 01293-422-911

Brazil ■ 0800-557779

Hong Kong ■ 800 933 965

The Netherlands ■ 0800 0229592

USA ■ 800-426-8157



Sample & Assay Technologies